



KCET Test–2022

PYQ

Maths

- The solution of the following equation  $\frac{dy}{dx} = (x+y)^2$  is
  - $\cot^{-1}(x+y) = x+c$
  - $\tan^{-1}(x+y) = x+c$
  - $\tan^{-1}(x+y) = 0$
  - $\cot^{-1}(x+y) = c$
- If  $y(x)$  be the solution of differential equation  $x \log x \frac{dy}{dx} + y = 2x \log x$ ,  $y(e)$  is equal to
  - $2e$
  - $e$
  - $0$
  - $2$
- If  $|\vec{a}| = 2$  and  $|\vec{b}| = 3$  and the angle between  $\vec{a}$  and  $\vec{b}$  is  $120^\circ$ , then the length of the vector  $\left| \frac{1}{2}\vec{a} - \frac{1}{3}\vec{b} \right|^2$  is
  - 1
  - 2
  - 3
  - $\frac{1}{6}$
- If  $|\vec{a} \times \vec{b}| + |\vec{a} \cdot \vec{b}|^2 = 36$  and  $|\vec{a}| = 3$  then  $|\vec{b}|$  is equal to
  - 2
  - 9
  - 36
  - 4
- If  $\vec{\alpha} = \hat{i} - 3\hat{j}$ ,  $\vec{\beta} = \hat{i} + 2\hat{j} - \hat{k}$  then express  $\vec{\beta}$  in the form  $\vec{\beta} = \vec{\beta}_1 + \vec{\beta}_2$  where  $\vec{\beta}_1$  is parallel to  $\vec{\alpha}$  and  $\vec{\beta}_2$  is perpendicular to  $\vec{\alpha}$  then  $\vec{\beta}_1$  is given by
  - $\hat{i} + 3\hat{j}$
  - $\frac{5}{8}(\hat{i} - 3\hat{j})$
  - $\frac{5}{8}(\hat{i} + 3\hat{j})$
  - $\hat{i} + 3\hat{j}$
- The sum of the degree and order of the differential equation  $(1+y_1^2)^{2/3} = y_2$  is
  - 7
  - 4
  - 6
  - 5
- If  $\frac{dy}{dx} + \frac{y}{x} = x^2$ , then  $2y(2) - y(1) =$ 
  - $\frac{13}{4}$
  - $\frac{11}{4}$
  - $\frac{15}{4}$
  - $\frac{9}{4}$
- The angle between the pair of lines  $\frac{x+3}{3} = \frac{y-1}{5} = \frac{z+3}{4}$  and  $\frac{x+1}{1} = \frac{y-4}{4} = \frac{z-5}{2}$  is
  - $\theta = \cos^{-1} \left[ \frac{5\sqrt{3}}{16} \right]$
  - $\theta = \cos^{-1} \left[ \frac{27}{5} \right]$
  - $\theta = \cos^{-1} \left[ \frac{8\sqrt{3}}{15} \right]$
  - $\theta = \cos^{-1} \left[ \frac{19}{21} \right]$
- The corner points of the feasible region of an LPP are  $(0,2), (3,0), (6,0), (6,8)$  and  $(0,5)$ , then the minimum value of  $z = 4x + 6y$  occurs at
  - Only two points
  - Finite number of points
  - Infinite number of points
  - Only one point



10. A dietician has to develop a special diet using two foods  $X$  and  $Y$ . Each packet (containing 30 g) of food.  $X$  contains 12 units of calcium, 4 units of iron, 6 units of cholesterol and 6 units of vitamin  $A$ . Each packet of the same quantity of food  $Y$  contains 3 units of calcium, 20 units of iron, 4 units of cholesterol and 3 units of vitamin  $A$ . The diet requires at least 240 units of calcium, at least 460 units iron and at most 300 units of cholesterol. The corner points of the feasible region are
- (A)  $(2, 72), (40, 15), (115, 0)$   
 (B)  $(2, 72), (40, 15), (15, 20)$   
 (C)  $(2, 72), (15, 20), (0, 23)$   
 (D)  $(0, 23), (40, 15), (2, 72)$
11. The distance of the point whose position vector is  $(2\hat{i} + \hat{j} - \hat{k})$  from the plane  $\vec{r} \cdot (\hat{i} - 2\hat{j} + 4\hat{k}) = 4$  is
- (A)  $\frac{-8}{21}$   
 (B)  $\frac{8}{\sqrt{21}}$   
 (C)  $8\sqrt{21}$   
 (D)  $\frac{-8}{\sqrt{21}}$
12. The co-ordination of foot of the perpendicular drawn from the origin to the plane  $2x - 3y + 4z = 29$  are
- (A)  $(-2, -3, 4)$       (B)  $(2, 3, 4)$   
 (C)  $(2, -3, -4)$       (D)  $(2, -3, 4)$
13. If  $A$  and  $B$  are two events such that  $P(A) = \frac{1}{2}, P(B) = \frac{1}{3}$  and  $P(A/B) = \frac{1}{4}$ , then  $P(A' \cap B')$  is
- (A)  $\frac{3}{4}$       (B)  $\frac{1}{4}$   
 (C)  $\frac{3}{16}$       (D)  $\frac{1}{12}$
14. A pandemic has been spreading all over the world. The probabilities are 0.7 that there will be a lockdown, 0.8 that the pandemic is controlled in one month if there is a lockdown and 0.3 that it is controlled in one month if there is no lockdown. The probability that the pandemic will be controlled in one month is
- (A) 0.46  
 (B) 0.65  
 (C) 1.65  
 (D) 1.46
15. If  $A$  and  $B$  are two independent events such that  $P(\bar{A}) = 0.75, P(A \cup B) = 0.65$ , and  $P(B) = x$ , then find the value of  $x$  :
- (A)  $\frac{7}{15}$   
 (B)  $\frac{5}{14}$   
 (C)  $\frac{8}{15}$   
 (D)  $\frac{9}{14}$
16. Find the mean number of heads in three tosses of a fair coin:
- (A) 3.5      (B) 1.5  
 (C) 4.5      (D) 2.5
17. The domain of the function  $f(x) = \frac{1}{\log_{10}(1-x)} + \sqrt{x+2}$  is
- (A)  $[-2, 0) \cup (0, 1)$   
 (B)  $[-2, 0) \cap (0, 1)$   
 (C)  $[-2, 1)$   
 (D)  $[-2, 0)$
18. The trigonometric function  $y = \tan x$  in the II quadrant
- (A) increases from  $-\infty$  to 0  
 (B) decreases from 0 to  $\infty$   
 (C) decreases from  $-\infty$  to 0  
 (D) increases from 0 to  $\infty$



19. The degree measure of  $\frac{\pi}{32}$  is equal to
- (A)  $4^{\circ}30'30''$   
 (B)  $5^{\circ}30'20''$   
 (C)  $5^{\circ}37'20''$   
 (D)  $5^{\circ}37'30''$
20. The value of  $\sin\frac{5\pi}{12}\sin\frac{\pi}{12}$  is
- (A)  $\frac{1}{4}$   
 (B) 0  
 (C) 1  
 (D)  $\frac{1}{2}$
21.  $\sqrt{2+\sqrt{2+\sqrt{2+2\cos 8\theta}}}$  =
- (A)  $2\cos\frac{\theta}{2}$   
 (B)  $\sin 2\theta$   
 (C)  $2\cos\theta$   
 (D)  $2\sin\theta$
22. If  $A = \{1, 2, 3, \dots, 10\}$  then number of subsets of  $A$  containing only odd numbers is
- (A) 30  
 (B) 31  
 (C) 27  
 (D) 32
23. Suppose that the number of elements in set  $A$  is  $p$ , the number of elements in set  $B$  is  $q$  and the number of elements in  $A \times B$  is 7 then  $p^2 + q^2 = \underline{\hspace{2cm}}$ .
- (A) 49 (B) 50  
 (C) 51 (D) 42
24. If  $a_1, a_2, a_3, \dots, a_{10}$  is a geometric progression and  $\frac{a_3}{a_1} = 25$ , then  $\frac{a_9}{a_5}$  equals
- (A)  $2(5^2)$  (B)  $3(5^2)$   
 (C)  $5^4$  (D)  $5^3$
25. If the straight line  $2x - 3y + 17 = 0$  is perpendicular to the line passing through the points  $(7, 17)$  and  $(15, \beta)$ , then  $\beta$  equals
- (A) -29  
 (B) -5  
 (C) 5  
 (D) 29
26. The octant in which the point  $(2, -4, -7)$  lies is
- (A) Fifth  
 (B) Eighth  
 (C) Third  
 (D) Fourth
27. If  $f(x) = \begin{cases} x^2 - 1, & 0 < x < 2 \\ 2x + 3, & 2 \leq x < 3 \end{cases}$ , the quadratic equation whose roots are  $\lim_{x \rightarrow 2^-} f(x)$  and  $\lim_{x \rightarrow 2^+} f(x)$  is
- (A)  $x^2 - 7x + 8 = 0$   
 (B)  $x^2 - 14x + 49 = 0$   
 (C)  $x^2 - 10x + 21 = 0$   
 (D)  $x^2 - 6x + 9 = 0$
28. If  $3x + i(4x - y) = 6 - i$  where  $x$  and  $y$  are real numbers, then the values of  $x$  and  $y$  are respectively,
- (A) 3, 4  
 (B) 3, 9  
 (C) 2, 4  
 (D) 2, 9
29. If all permutations of the letters of the word MASK are arranged in the order as in dictionary with or without meaning, which one of the following is 19<sup>th</sup> word?
- (A) AMSK  
 (B) KAMS  
 (C) SAMK  
 (D) AKMS



30. If the set  $x$  contains 7 elements and set  $y$  contains 8 elements, then the number of bijections from  $x$  to  $y$  is  
 (A)  $8!$   
 (B)  $0$   
 (C)  $8P_7$   
 (D)  $7!$
31. If  $f : R \rightarrow R$  be defined  $f(x) = \begin{cases} 2x & : x > 3 \\ x^2 & : 1 < x \leq 3 \\ 3x & : x \leq 1 \end{cases}$   
 Then  $f(-1) + f(2) + f(4)$  is  
 (A)  $14$   
 (B)  $5$   
 (C)  $10$   
 (D)  $9$
32. Let the relation  $R$  is defined in  $N$  by a  $Rb$ , if  $3a + 2b = 27$  then  $R$  is  
 (A)  $\{(2,1)(9,3)(6,5)(3,7)\}$   
 (B)  $\{(1,12)(3,9)(5,6)(7,3)\}$   
 (C)  $\left\{ \left( 0, \frac{27}{2} \right) (1,12)(3,9)(5,6)(7,3) \right\}$   
 (D)  $\{(1,12)(3,9)(5,6)(7,3)(9,0)\}$
32.  $\lim_{y \rightarrow 0} \frac{\sqrt{3+y^3} - \sqrt{3}}{y^3} =$   
 (A)  $3\sqrt{2}$   
 (B)  $\frac{1}{2\sqrt{3}}$   
 (C)  $\frac{1}{3\sqrt{2}}$   
 (D)  $2\sqrt{3}$
34. If the standard deviation of the numbers  $-1, 0, 1, k$  is  $\sqrt{5}$  where  $k > 0$ , then  $k$  is equal to  
 (A)  $2\sqrt{6}$                       (B)  $4\sqrt{\frac{5}{3}}$   
 (C)  $\sqrt{6}$                          (D)  $2\sqrt{\frac{10}{3}}$
35. If  $A = \begin{bmatrix} 2 & -1 \\ 3 & -2 \end{bmatrix}$ , then the inverse of the matrix  $A^3$  is  
 (A)  $-A$   
 (B)  $A$   
 (C)  $-1$   
 (D)  $1$
36. If  $A$  is a skew symmetric matrix, then  $A^{2021}$  is  
 (A) Skew symmetric matrix  
 (B) Row matrix  
 (C) Column matrix  
 (D) Symmetric matrix
37. If  $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$  then  $(aI + bA)^n$  is (where  $I$  is the identity matrix of order 2)  
 (A)  $a^n I + b^n A$   
 (B)  $a^2 I + a^{n-1} b \cdot A$   
 (C)  $a^n I + n \cdot a^{n-1} b \cdot A$   
 (D)  $a^n I + na^n bA$
38. If  $A$  is a  $3 \times 3$  matrix such that  $|5 \cdot \text{adj}A| = 5$  then  $|A|$  is equal to  
 (A)  $\pm 5$   
 (B)  $\pm 1$   
 (C)  $\pm 1/25$   
 (D)  $\pm 1/5$
39. If there are two values of '  $a$  ' which makes determinant  $\Delta = \begin{vmatrix} 1 & -2 & 5 \\ 2 & a & -1 \\ 0 & 4 & 2a \end{vmatrix}$  Then the sum of these numbers is  
 (A)  $5$   
 (B)  $-4$   
 (C)  $9$   
 (D)  $4$



40. If the vertices of a triangle are  $(-2,6)$ ,  $(3,-6)$  and  $(1,5)$ , then the area of the triangle is  
 (A) 35sq.units  
 (B) 40 sq. Units  
 (C) 15. 5sq.units  
 (D) 30 sq. Units
41. Domain of  $\cos^{-1}[x]$  is, where  $[ ]$  denotes a greatest integer function  
 (A)  $[-1,2)$   
 (B)  $(-1,2]$   
 (C)  $(-1,2)$   
 (D)  $[-1,2]$
42. If  $A$  is matrix of order  $3 \times 3$ , then  $(A^2)^{-1}$  is equal to  
 (A)  $(-A)^{-2}$   
 (B)  $(-A^2)^2$   
 (C)  $(A^{-1})^2$   
 (D)  $A^2$
43. If  $x = e^\theta \sin \theta$ ,  $y = e^\theta \cos \theta$  where  $\theta$  is a parameter, then  $\frac{dy}{dx}$  at  $(1,1)$  is equal to  
 (A)  $-\frac{1}{4}$   
 (B) 0  
 (C)  $\frac{1}{2}$   
 (D)  $-\frac{1}{2}$
44. If  $y = e^{\sqrt{x}\sqrt{x}\sqrt{x}\dots}$ ,  $x > 1$  then  $\frac{d^2y}{dx^2}$  at  $x = \log_e^3$  is  
 (A) 1  
 (B) 3  
 (C) 5  
 (D) 0
45. If  $f(1)=1, f'(1)=3$  then the derivative of  $f(f(f(x))) + (f(x))^2$  at  $x=1$  is  
 (A) 12  
 (B) 10  
 (C) 33  
 (D) 35
46. If  $y = x^{\sin x} + (\sin x)^x$  then  $\frac{dy}{dx}$  at  $x = \frac{\pi}{2}$  is  
 (A)  $\frac{\pi^2}{2}$   
 (B)  $\frac{4}{\pi}$   
 (C)  $\pi \log \frac{\pi}{2}$   
 (D) 1
47. If  $A_n = \begin{bmatrix} 1-n & n \\ n & 1-n \end{bmatrix}$  then  $|A_1| + |A_2| + \dots + |A_{2021}| =$   
 (A) 4042  
 (B) -2021  
 (C)  $-(2021)^2$   
 (D)  $(2021)^2$
48. If  $y = (1+x^2)\tan^{-1}x - x$  then  $\frac{dy}{dx}$  is  
 (A)  $x\tan^{-1}x$   
 (B)  $2x\tan^{-1}x$   
 (C)  $\frac{\tan^{-1}x}{x}$   
 (D)  $x^2\tan^{-1}x$
49. The co-ordinates of the point on the  $\sqrt{x} + \sqrt{y} = 6$  at which the tangent is equally inclined to the axes is  
 (A) (6,6)  
 (B) (4,4)  
 (C) (1,1)  
 (D) (9,9)



50. The function  $f(x) = 4\sin^3 x - 6\sin^2 x + 12\sin x + 100$  is strictly
- (A) decreasing in  $\left(\frac{\pi}{2}, \pi\right)$
- (B) decreasing in  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
- (C) decreasing in  $\left[0, \frac{\pi}{2}\right]$
- (D) increasing in  $\left(\pi, \frac{3\pi}{2}\right)$
51. If  $[x]$  is the greatest integer function not greater than  $x$  then  $\int_0^8 [x] dx$  is equal to
- (A) 20
- (B) 28
- (C) 30
- (D) 29
52.  $\int_0^{\pi/2} \sqrt{\sin \theta} \cos^3 \theta d\theta$  is equal to
- (A)  $\frac{7}{21}$
- (B)  $\frac{8}{23}$
- (C)  $\frac{7}{23}$
- (D)  $\frac{8}{21}$
53. If  $e^y + xy = e$  the order pair  $\left(\frac{dy}{dx}, \frac{d^2y}{dx^2}\right)$  at  $x = 0$  is equal to
- (A)  $\left(\frac{-1}{e}, \frac{1}{e^2}\right)$
- (B)  $\left(\frac{1}{e}, \frac{1}{e^2}\right)$
- (C)  $\left(\frac{-1}{e}, \frac{-1}{e^2}\right)$
- (D)  $\left(\frac{1}{e}, \frac{-1}{e^2}\right)$
54. The function  $f(x) = \log(1+x) - \frac{2x}{2+x}$  is increasing on
- (A)  $(-\infty, 0)$
- (B)  $(-\infty, \infty)$
- (C)  $(\infty, -1)$
- (D)  $(-1, \infty)$
55.  $\int_0^1 \frac{xe^x}{(2+x)^3} dx$  is equal to
- (A)  $\frac{1}{9} \cdot e - \frac{1}{4}$
- (B)  $\frac{1}{27} \cdot e - \frac{1}{8}$
- (C)  $\frac{1}{27} \cdot e + \frac{1}{8}$
- (D)  $\frac{1}{9} \cdot e + \frac{1}{4}$
56. If  $\int \frac{dx}{(x+2)(x^2+1)} = a \log|1+x^2| + b \tan^{-1} x + \frac{1}{5} \log|x+2| + c$  then
- (A)  $a = \frac{1}{10} b = \frac{-2}{5}$
- (B)  $a = \frac{-1}{10} b = \frac{2}{5}$
- (C)  $a = \frac{1}{10} b = \frac{2}{5}$
- (D)  $a = \frac{-1}{10} b = \frac{-2}{5}$
57. Area of the region bounded by the curve  $y = \tan x$ , the  $x$ -axis and the line  $x = \frac{\pi}{3}$  is
- (A)  $-\log 2$
- (B)  $\log \frac{1}{2}$
- (C)  $\log 2$
- (D) 0



58. Evaluate  $\int_2^3 x^2 dx$  as the limit of a sum

(A)  $\frac{19}{3}$                       (B)  $\frac{72}{6}$

(C)  $\frac{53}{9}$                       (D)  $\frac{25}{7}$

59.  $\int_0^{\pi/2} \frac{\cos x \sin x}{1 + \sin x} dx$  is equal to

(A)  $1 - \log 2$                 (B)  $\log 2 - 1$

(C)  $\log 2$                       (D)  $-\log 2$

60.  $\int \frac{\cos 2x - \cos 2\alpha}{\cos x - \cos \alpha} dx$  is equal to

(A)  $2(\sin x + 2x \cos \alpha) + c$

(B)  $2(\sin x - x \cos \alpha) + c$

(C)  $2(\sin x + x \cos \alpha) + c$

(D)  $2(\sin x - 2x \cos \alpha) + c$