



KCET Test–2025

PYQ

Chemistry

1. Match List-I with List-II and select the correct option:

List-I (Molecule/ion)		List-II (Bond order)	
1.	NO	i.	1.5
2.	CO	ii.	2.0
3.	O ₂	iii.	2.5
4.	O ₂	iv.	3.0

- (A) a-iii, b-iv, c-i, d-ii
 (B) a-i, b-iv, c-iii, d-ii
 (C) a-ii, b-iii, c-iv, d-i
 (D) a-iv, b-iii, c-ii, d-i
2. The electronic configuration of X and Y are given below:
 X: $1s^2 2s^2 2p^6 3s^2 3p^3$
 Y: $1s^2 2s^2 2p^6 3s^2 3p^5$
 Which of the following is the correct molecular formula and type of bond formed between X and Y?
 (A) X₃Y, ionic bond
 (B) X₂Y₃, coordinate bond
 (C) XY₃, covalent bond
 (D) X₂Y, covalent bond

3. Match List-I with List-II

List-I (Types of redox reactions)		List-II (Examples)	
a.	Combination reaction	i.	$\text{Cl}_{2(g)} + 2\text{Br}^-_{(aq)} \rightarrow 2\text{Cl}^-_{(aq)} + \text{Br}_{2(l)}$
b.	Decomposition reaction	ii.	$2\text{H}_2\text{O}_{2(aq)} \rightarrow 2\text{H}_2\text{O}_{(l)} + \text{O}_{2(g)}$
c.	Displacement reaction	iii.	$\text{CH}_{4(g)} + 2\text{O}_{2(g)} \xrightarrow{\Delta} \text{CO}_{2(g)} + 2\text{H}_2\text{O}_{(l)}$
d.	Disproportionation reaction	iv.	$2\text{H}_2\text{O}_{(l)} \xrightarrow{\Delta} 2\text{H}_{2(g)} + \text{O}_{2(g)}$

Choose the correct answer from the options given below:

- (A) a-iv, b-iii, c-i, d-ii
 (B) a-ii, b-i, c-iv, d-iii
 (C) a-iii, b-iv, c-i, d-ii
 (D) a-iii, b-ii, c-i, d-iv

4. In the following pairs, the one in which both transition metal ions are colourless is
 (A) $\text{Sc}^{3+}, \text{Zn}^{2+}$ (B) $\text{V}^{2+}, \text{Ti}^{3+}$
 (C) $\text{Zn}^{2+}, \text{Mn}^{2+}$ (D) $\text{Ti}^{4+}, \text{Cu}^{2+}$
5. In the reaction between hydrogen sulphide and acidified permanganate solution,
 (A) H₂S is reduced to S, MnO_4^- is oxidised to Mn^{2+}
 (B) H₂S is oxidised to SO₂, MnO_4^- is reduced to MnO_2
 (C) H₂S is reduced to SO₂, MnO_4^- is oxidised to Mn^{2+}
 (D) H₂S is oxidised to S, MnO_4^- is reduced to Mn^{2+}
6. A member of the Lanthanoid series which is well known to exhibit +4 oxidation state is
 (A) Samarium (B) Europium
 (C) Erbium (D) Cerium
7. In which of the following pairs, both the elements do not have $(n-1)d^{10}ns^2$ configuration?
 (A) Cu, Zn (B) Zn, Cd
 (C) Cd, Hg (D) Ag, Cu
8. A ligand which has two different donor atoms and either of the two ligates with the central metal atom/ion in the complex is called_____
 (A) Chelate ligand
 (B) Unidentate ligand
 (C) Polydentate ligand
 (D) Ambidentate ligand
9. Which of the following statements are true about $[\text{NiCl}_4]^{2-}$?
 (a) The complex has tetrahedral geometry
 (b) Co-ordination number of Ni is 2 and oxidation state is +4
 (c) The complex is sp^3 hybridised
 (d) It is a high spin complex
 (e) The complex is paramagnetic
 (A) a, c, d and e (B) a, b, d and e
 (C) b, c, d and e (D) a, b, c and d



10. Which formula and its name combination is incorrect?

- (A) $K_3[Cr(C_2O_4)_3]$, Potassium trioxalatochromate (III)
 (B) $[CoCl_2(en)_2]Cl$, Dichloridobis (ethane-1, 2-dimine) cobalt (III) chloride
 (C) $[Co(NH_3)_5(CO_3)]Cl$, Pentaamine carbonylcobalt (III) chloride
 (D) $[Pt(NH_3)_2Cl(NO_2)]$ Diamine chloridonitrito – N – Platinum (II)

11. In the complex ion $[Fe(C_2O_4)_3]^{3-}$, the coordination number of Fe is

- (A) 4 (B) 5
 (C) 6 (D) 3

12. Match List-I with List-II for the following reaction pattern

Glucose $\xrightarrow{\text{Reagent}}$ Product \longrightarrow Structural prediction

List-I (Reagents)		List-II (Structural prediction)	
a.	Acetic anhydride	i.	Glucose has an aldehyde group
b.	Bromine water	ii.	Glucose has a straight chain of six carbon atoms
c.	Hydroiodic acid	iii.	Glucose has five hydroxyl group
d.	Hydrogen cyanide	iv.	Glucose has a carbonyl group

Choose the correct answer from the options given below:

- (A) a-iv, b-iii, c-ii, d-i
 (B) a-iii, b-i, c-ii, d-iv
 (C) a-i, b-ii, c-iii, d-iv
 (D) a-iii, b-ii, c-i, d-iv

13. The correct sequence of α -amino acids, hormone, vitamin, carbohydrates respectively is

- (A) Thiamine, Thyroxine, Vitamin A, Glucose
 (B) Glutamine, Insulin, Aspartic acid, Fructose
 (C) Arginine, Testosterone, Glutamic acid, Fructose
 (D) Aspartic acid, Insulin, Ascorbic acid, rhamnose

14. Which examples of carbohydrates exhibit α -link, (α -glycosidic link) in their structure?

- (A) Maltose and Lactose

(B) Amylose and Amylopectin

(C) Cellulose and Glycogen

(D) Glucose and Fructose

15. In the titration of potassium permanganate ($KMnO_4$) against Ferrous ammonium sulphate (FAS) solution, dilute sulphuric acid but not nitric acid is used to maintain acidic medium, because

- (A) It is difficult to identify the end point
 (B) Nitric acid doesn't act as an indicator
 (C) Nitric acid itself is an oxidizing agent
 (D) Nitric acid is a weak acid than sulphuric acid

16. The group reagent $NH_4Cl(s)$ and aqueous NH_3 will precipitate which of the following ion?

- (A) NH_4^+ (B) Al^{3+}
 (C) Ba^{2+} (D) Ca^{2+}


17. In the preparation of sodium fusion extract, the purpose of fusing organic compound with a piece of sodium metal is to


- (A) Convert the organic compound into vapour state
 (B) Convert the elements of the compound from covalent form to ionic form
 (C) Convert the elements of the compound from ionic form to covalent form
 (D) Decrease the melting point of the compound

18. The sodium fusion extract is boiled with concentrated nitric acid while testing for halogens. By doing so, it


- (A) helps in precipitation of $AgCl$
 (B) increases the solubility of $AgCl$
 (C) increases the concentration of NO_3^- ion
 (D) decomposes Na_2S and $NaCN$, if formed

19. Which of the following is not an aromatic compound






20. The IUPAC name of the given organic compound is




- (A) Hexa-1-yn-3,5-diene
 (B) Hexa-5-yn-1,3-diene
 (C) Hexa-1,3-dien-5-yne
 (D) Hexa-3,5-dien-1-yne


21. Among the following, identify the compound that is not an isomer of hexane




(A)




(B)



(C)



(D)

22. The organic compound  can be

classified as _____

- (A) Allylic halide (B) Benzyl halide
 (C) Aryl halide (D) Alkyl halide


23. Chlorobenzene reacts with bromine gas in the presence of Anhyd AlBr_3 to yield p-Bromochlorobenzene. This reaction is classified as _____

- (A) Elimination reaction
 (B) Nucleophilic substitution reaction
 (C) Electrophilic substitution reaction
 (D) Addition reaction

24. The organometallic compound $(\text{CH}_3)_3 \text{CMgBr}$ on reaction with D_2O produces _____

- (A) $(\text{CH}_3)_3\text{COD}$ (B) $(\text{CD}_3)_3\text{CD}$
 (C) $(\text{CD}_3)_3\text{COD}$ (D) $(\text{CH}_3)_3\text{CD}$

25. The major product formed when 1-Bromo-3-Chlorocyclobutane reacts with metallic sodium in dry ether is



26. Ethyl alcohol is heated with concentrated sulphuric acid at 413 K. The major product

- (A) $\text{C}_2\text{H}_5 - \text{O} - \text{C}_2\text{H}_5$ (B) $\text{CH}_3 - \text{O} - \text{C}_3\text{H}_7$
 (C) $\text{CH}_2 = \text{CH}_2$ (D) $\text{CH}_3\text{COOC}_2\text{H}_5$

27. Phenol can be distinguished from propanol by using the reagent


- (A) Bromine water (B) Iron metal
 (C) Iodine in alcohol (D) Sodium metal







28. Match the following with their pKa values

Acid	pKa
(I) Phenol	(a) 16
(II) p-Nitrophenol	(b) 0.78
(III) Ethyl alcohol	(c) 10
(IV) Picric acid	(d) 7.1

- (A) I-c, II-d, III-a, IV-b
 (B) I-a, II-d, III-c, IV-b
 (C) I-a, II-b, III-c, IV-d
 (D) I-b, II-a, III-d, IV-c



A and B respectively are



30. Oxidation of Toluene with chromyl chloride followed by hydrolysis gives Benzaldehyde. This reaction is known as _____

- (A) Etard Reaction (B) Kolbe reaction
 (C) Stephen reaction (D) Cannizzaro Reaction

31. Statement – I : Reduction of ester by DIABL-H followed by hydrolysis gives aldehyde. Statement – II : Oxidation of benzyl alcohol with aqueous KMnO_4 leads to the formation to Benzaldehyde.

Among the above statements, identify the correct statement.

- (A) Both statements – I and II are false

- (B) Statement – I is true but statement – II is false
 (C) Statement – I is false but statement – II is true
 (D) Both statements – I and II are true.

32. Arrange the following compounds in their decreasing order of reactivity towards Nucleop addition reaction.



- (A) $\text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{COC}_2\text{H}_5$
 (B) $\text{CH}_3\text{COCH}_3 > \text{CH}_3\text{CHO} > \text{CH}_3\text{COC}_2\text{H}_5$
 (C) $\text{CH}_3\text{COC}_2\text{H}_5 > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{CHO}$
 (D) $\text{CH}_3\text{CHO} > \text{CH}_3\text{COC}_2\text{H}_5 > \text{CH}_3\text{COCH}_3$





33. Which of the following has most acidic Hydrogen?

- (A) Propanoic acid
 (B) Dichloroacetic acid
 (C) Trichloroacetic acid
 (D) Chloroacetic acid

34. Which of the following reagents are suitable to differentiate Aniline and N-methylaniline chemical

- (A) Acetic anhydride
 (B) Br_2 water
 (C) Conc. Hydrochloric acid and anhydrous zinc chloride
 (D) Chloroform and Alcoholic potassium hydroxide

35. Which of the following reaction/s does not yield an amine ?







- (A) Both I and III (B) Only II
 (C) Only III (D) Both II and IV



36. Match the compounds given in List – I with the items given in List – II.

List-I		List-II	
(I)	Benzenesulphonyl Chloride	(a)	Zwitterioin
(II)	Sulphanilic acid	(b)	Hinsberg reagent
(III)	Alkyl Diazonium salts	(c)	Dyes
(IV)	Aryl Diazonium salts	(d)	Conversion to alcohols

- (A) I-c, II-b, III-a, IV-d
 (B) I-a, II-c, III-b, IV-d
 (C) I-c, II-a, III-d, IV-b
 (D) I-b, II-a, III-d, IV-c
37. The number of orbitals associated with 'N' shell of an atom is
 (A) 16 (B) 32
 (C) 3 (D) 4
38. According to the Heisenberg's Uncertainty principle, the value of $\Delta b \cdot \Delta x$ for an object whose mass is 10^{-6} kg is ($h = 6.626 \times 10^{-34}$ Js)
 (A) $3.0 \times 10^{-24} \text{ m}^{-2}\text{s}^{-1}$
 (B) $4.0 \times 10^{-26} \text{ m}^{-2}\text{s}^{-1}$
 (C) $3.5 \times 10^{-25} \text{ m}^{-2}\text{s}^{-1}$
 (D) $5.2 \times 10^{-29} \text{ m}^{-2}\text{s}^{-1}$
39. Given below are two statements.
Statement-I : Adiabatic work done is positive when work is done on the system and internal energy of the system increases.
Statement – II : No work is done during free expansion of an ideal gas.
 In the light of the above statements, choose the correct answer from the options given below.
 (A) Both statements – I and Statement – II are false
 (B) Statement – I is true but statement – II is false
 (C) Statement – I is false but statement – II is true
 (D) Both statements – I and Statement – II are true.
40. Which one of the following reactions has $\Delta H = \Delta U$?




41. Identify the incorrect statements among the following:
 (a) All enthalpies of fusion are positive
 (b) The magnitude of enthalpy change does not depend on the strength of the intermolecular interactions in the substance undergoing phase transformations.
 (c) When a chemical reaction is reversed, the value of $\Delta_r H^\circ$ is reversed in sign.
 (d) The change in enthalpy is dependent of path between initial state (reactants) and final state (products)
 (e) For most of the ionic compounds, $\Delta_{\text{sol}} H^\circ$ is negative
 (A) a, b and d (B) b, d and e
 (C) a, d and e (D) a and e only
42. Which of the following statements is/are true about equilibrium?
 (a) Equilibrium is possible only in a closed system of at a given temperature
 (b) All the measurable properties of the system remain constant at equilibrium
 (c) Equilibrium constant for the reverse reaction is the inverse of the equilibrium constant for the reaction in the forward direction.
 (A) Only b (B) Only c
 (C) a, b and c (D) Only a
43. According to Le Chatelier's principle, in the reaction $\text{CO}(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g})$, the formation of methane is favoured by
 (a) Increasing the concentration of CO
 (b) Increasing the concentration of H_2O
 (c) Decreasing the concentration of CH_4
 (d) Decreasing the concentration of H_2
 (A) a and c (B) b and d
 (C) a and d (D) a and b



44. The equilibrium constant at 298K for the reaction $A + B \rightleftharpoons C + D$ is 100. If the initial concentrations of all the four species were 1M each, then equilibrium concentration of D (in mol L⁻¹) will be
- (A) 0.182 (B) 1.818
(C) 1.182 (D) 0.818

45. Among the following 0.1 m aqueous solutions, which one will exhibit the lowest boiling point elevation, assuming complete ionization of the compound in solution?
- (A) Aluminium chloride
(B) Aluminium sulphate
(C) Potassium sulphate
(D) Sodium chloride

46. Variation of solubility with temperature t for a gas in liquid is shown by the following graphs. The correct representation is



47. 180g of glucose, C₆H₁₂O₆, is dissolved in 1 kg of water in a vessel. The temperature at which water boils at 1.013 bar is ____ (given, K_b for water is 052K kg mol⁻¹. Boiling point for pure water is 373.15K)
- (A) 373.67 K (B) 373.015 K
(C) 373.0 K (D) 373.202 K

48. If N₂ gas is bubbled through water at 293 K, how many moles of N₂ gas would dissolve in 1 litre of water? Assume that N₂ exerts a partial pressure of 0.987 bar.

[Given K_H for N₂ at 293 K is 76.48 K bar]

- (A) 0.716×10^{-3} (B) 7.16×10^{-5}
(C) 7.16×10^{-4} (D) 7.16×10^{-3}
49. The correct statement/s about Galvanic cell is/are
- (a) Current flows from cathode to anode
(b) Anode is positive terminal
(c) If cell E_{cell} < 0 then it is spontaneous reaction
(d) Cathode is positive terminal
- (A) a and b only (B) a, b and c
(C) a and d only (D) b only
50. The electronic conductance depends on
- (A) Nature of electrolyte added
(B) The number of valence electrons per atom
(C) Concentration of the electrolyte
(D) Size of the ions
51. For a given half cell, $Al^{3+} + 3e^{-} \rightarrow Al$ on increasing of aluminium ion, the electrode potential will
- (A) Decrease
(B) No change
(C) First increase then decrease
(D) Increase

52. Match the following select the correct option for the quantity of electricity, in Cmol⁻¹ required to deposit various metals at cathode

List-I		List-II	
a.	Ag ⁺	i.	386000 Cmol ⁻¹
b.	Mg ²⁺	ii.	289500 Cmol ⁻¹
c.	Al ³⁺	iii.	96500Cmol ⁻¹
d.	Ti ⁴⁺	iv.	193000 Cmol ⁻¹


- (A) a-ii, b-I, c-iv, d-iii
(B) a-iii, b-iv, c-ii, d-i
(C) a-iv, b-iii, c-i, d-ii
(D) a-I, b-ii, c-iii, d-iv



53. Catalysts are used to increase the rate of a chemical reaction. Because it
- (A) Increases the activation energy of the reaction
 (B) Decreases the activation energy of the reaction
 (C) Brings about improper orientation of reactant molecules
 (D) Increases the potential energy barrier

54. Half-life of a first order reaction is 20 seconds and initial concentration of reactant is 0.2M. The concentration of reactant left after 80 seconds is
- (A) 0.1 M (B) 0.05 M
 (C) 0.0125 M (D) 0.2 M

55. In the given graph, a E for the reverse reaction will be



- (A) 125 KJ (B) 215 KJ
 (C) 90 KJ (D) 305 KJ
56. For the reaction $2\text{N}_2\text{O}_{5(g)} \rightarrow 4\text{NO}_{2(g)} + \text{O}_{2(g)}$ initial concentration of N_2O_5 is 2.0 mol L^{-1} and after 300 min, it is reduced to 1.4 mol L^{-1} . The rate of production of NO_2 (in $\text{mol L}^{-1} \text{ min}^{-1}$) is
- (A) 2.5×10^{-4} (B) 4×10^{-4}
 (C) 2.5×10^{-3} (D) 4×10^{-3}
57. Which of the following methods of expressing concentration are unitless?
- (A) Mole fraction and Mass percent (W/W)

- (B) Molality and Mole fraction
 (C) Mass percent (W/W) and Molality
 (D) Molality and Molarity

58. Select the INCORRECT statement/s from the following:
- (a) 22 books have infinite significant figures
 (b) In the answer of calculation 2.5×1.25 has four significant figures,
 (c) Zero's preceding to first non-zero digit are significant
 (d) In the answer of calculation $12.11 + 18.0 + 1.012$ has three significant figures
- (A) b, c and d (B) b and c only
 (C) b and d only (D) a and b only

59. Given below and the atomic masses of the elements:

Element	Li	Na	Cl	K	Ca	Br	Sr	I	Ba
Atomic Mass (g mol ⁻¹):	7	23	35.5	39	40	80	88	127	137

Which of the following doesn't form triad?

- (A) Ba, Sr, Ca (B) Cl, Br, I
 (C) Cl, K, Ca (D) Li, Na, K
60. The change in hybridization (if any) of the 'Al' atom in the following reaction is
- $$\text{AlCl}_3 + \text{Cl}^- \longrightarrow \text{AlCl}_4^-$$
- (A) No change in the hybridization state
 (B) sp^2 to sp^3
 (C) sp^3 to sp^3d
 (D) sp^3 to sp^2