



PRACHAND NEET



ONE SHOT



Zoology

Biotechnology Principles & Processes

3-5
Ques

Dr. Akanksha Agarwal Ma'am



Topics *to be covered*

1

Biotechnology Principles & Processes
Complete chapter



You can't go back and
change the beginning,
but you can start
where you are and
change the ending.

—C.S. LEWIS



3 tips for
last 3 months

1. Regularity
2. Positivity
3. listen to yr self

PRACHAND SERIES

TELEGRAM CHANNEL



@PW_YAKEENDROPPER

Biotechnology deals with techniques of using live organisms or enzymes from organisms to produce products and processes useful to humans. In this sense, making curd, bread or wine, which are all microbe-mediated processes, could also be thought as a form of biotechnology. However, it is used in a restricted sense today, to refer to such of those processes which use genetically modified organisms to achieve the same on a larger scale. Further, many other processes/techniques are also included under biotechnology. For example, *in vitro* fertilisation leading to a 'test-tube' baby, synthesising a gene and using it, developing a DNA vaccine or correcting a defective gene, are all part of biotechnology.

The European Federation of Biotechnology (EFB) has given a definition of biotechnology that encompasses both traditional view and modern molecular biotechnology.

The definition given by EFB is as follows:

'The integration of natural science and organisms, cells, parts thereof, and molecular analogues for products and services'.

9.1 PRINCIPLES OF BIOTECHNOLOGY

Among many, the two core techniques that enabled birth of modern biotechnology are :

- (i) **Genetic engineering** : Techniques to ^{change} alter the chemistry of genetic material (DNA and RNA),

to introduce these into host organisms and thus change the phenotype of the host organism.

- (iii) **Bioprocess engineering**: Maintenance of sterile (microbial contamination-free) ambience in chemical engineering processes to enable growth of only the desired microbe/eukaryotic cell in large quantities for the manufacture of biotechnological products like antibiotics, vaccines, enzymes, etc.

Let us now understand the conceptual development of the principles of genetic engineering.

You probably appreciate the advantages of sexual reproduction over asexual reproduction. The former provides opportunities for variations and formulation of unique combinations of genetic setup, some of which may be beneficial to the organism as well as the population. Asexual reproduction preserves the genetic information, while sexual reproduction permits variation. Traditional hybridisation procedures used in plant and animal breeding, very often lead to inclusion and multiplication of

undesirable genes along with the desired genes. The techniques of genetic engineering which include creation of **recombinant DNA**, use of **gene cloning** and **gene transfer**, overcome this limitation and allows us to isolate and introduce only one or a set of desirable genes without introducing undesirable genes into the target organism.

Do you know the likely fate of a piece of DNA, which is somehow transferred into an alien organism? Most likely, this piece of DNA would not be able to multiply itself in the progeny cells of the organism. But, when it gets integrated into the genome of the recipient, it may multiply and be inherited along with the host DNA. This is because the alien piece of DNA has become part of a chromosome, which has the ability to replicate. In a chromosome there is a specific DNA sequence called the **origin of replication**, which is responsible for initiating replication. Therefore, for the multiplication of any alien piece of DNA in an organism

* Both cut with Same RE

* Desired gene + Vector DNA with ori (origin of Replication)

Foreign
Alien

Gene of Interest (GOI)

DNA LIGASE



HOST eg: E. coli
bacteria
Binary fission



III Gene Cloning

it needs to be a part of a chromosome(s) which has a specific sequence known as 'origin of replication'. Thus, an alien DNA is linked with the origin of replication, so that, this alien piece of DNA can replicate and multiply itself in the host organism. This can also be called as **cloning** or making multiple identical copies of any template DNA.


Let us now focus on the first instance of the construction of an artificial recombinant DNA molecule. The construction of the first recombinant DNA emerged from the possibility of linking a gene encoding antibiotic resistance with a native **plasmid** (autonomously replicating circular extra-chromosomal DNA) of *Salmonella typhimurium*. Stanley Cohen and Herbert Boyer accomplished this in 1972 by isolating the antibiotic resistance gene by cutting out a piece of DNA from a plasmid which was responsible for conferring antibiotic resistance. The cutting of DNA at specific locations became possible with the discovery of the so-called

~~V. amn~~ *

Construction of 1st rDNA: 1972, Stanley Cohen and Herbert Boyer



* Salmonella typhimurium



Small Circular, ds DNA, extra-chromosomal self/Autonomously replicating

Oxi ⊕

RE



* Desired gene (Antibiotic resist. gene)


DNA ligase

I rDNA

II Gene transfer



III Gene cloning



* Genetically Modified Organism
 Gmo
Transgenic organism
 HOST

} E. coli

'molecular scissors'– **restriction enzymes**. The cut piece of DNA was then linked with the plasmid DNA. These plasmid DNA act as **vectors** to transfer the piece of DNA attached to it. You probably know that mosquito acts as an insect vector to transfer the malarial parasite into human body. In the same way, a plasmid can be used as vector to deliver an alien piece of DNA into the host organism. The linking of antibiotic resistance gene with the plasmid vector became possible with the enzyme DNA ligase, which acts on cut DNA molecules and joins their ends. This makes a new combination of circular autonomously replicating DNA created *in vitro* and is known as recombinant DNA. When this DNA is transferred into *Escherichia coli*, a bacterium closely related to *Salmonella*, it could replicate using the new host's DNA polymerase enzyme and make multiple copies. The ability to multiply copies of antibiotic resistance gene in *E. coli* was called **cloning** of antibiotic resistance gene in *E. coli*.

You can hence infer that there are three basic steps in genetically modifying an organism —

- (i) identification of DNA with desirable genes;
- (ii) introduction of the identified DNA into the host;
- (iii) maintenance of introduced DNA in the host and transfer of the DNA to its progeny.

9.2 TOOLS OF RECOMBINANT DNA TECHNOLOGY

Now we know from the foregoing discussion that genetic engineering or recombinant DNA technology can be accomplished only if we have the key tools, i.e., restriction enzymes, polymerase enzymes, ligases, vectors and the host organism. Let us try to understand some of these in detail.



I ENZYMES :-

1. LYASES — To break open Cell and Isolation of purified DNA / Genetic mat.

- Digest Cell wall

- a) Cellulase, Pectinase (Plant / Algal cell)

- b) Chitinase (Fungal cell)

- c) Lysozyme (Bacterial cell, digest Peptidoglycan)

- Digest Plasma mb & Nuclear mb : Detergents (Lipase)

- Digest RNA —————→ RNAase (Ribonuclease enzyme)

- Digest Proteins —————→ Protease



P40 ↓ *
Addition of Chilled Ethanol

↓
DNA Precipitate out as
Fine threads in suspension




↓
Isolated Purified DNA
Removed by Spooling * P40



2. Restriction Endonuclease / Restriction Enzyme To cut ^(RE) DNA into fragments Molecular Scissors :-

NUCLEASE

[digest N.A
break 3,5-
phosphodiester]



(b) **Endonuclease**

- Cut in between in one strand only

(c) **RE**

- Inspect whole DNA & cut DNA at specific Palindromic / restriction / recognition site in Both strands



* Palindromic DNA which reads same on both strands if orientation of reading kept same

MADAM


MALAYALAM

SIMSI

STETJ



* Discovery of RE :- In 1963, Bacteria has 2 Enzyme complex



* One Cuts DNA
(viral)
called RE

* Bacteria's Natural
Defence mechanism,
to protect from
Bacteriophage infection

* other add
Methyl group
to bacteria's
own DNA

↓
Prevent bact.
suicide




* 1968, 1st RE isolated


* (5 yr. after discovery)

Hind II

recognise
6 bp long nucleotide
sequence
NCERT

* > 900 RE isolated
from 230 strains of Bacteria






Hind II

↓ ↓ ↓

Genus Species Strain
Haemophilus influenzae Rd



* Working of RE : Cut DNA on Both strands,
a little away from Centre of
Palindrome



✓ Imp * Sticky ends form H-bonds with complementary counterparts

↓
This Stickiness facilitates action of DNA LIGASE Enzyme (Phosp. diester bond formation)
PYQ

↓
rDNA formed

* Both Vector DNA & Foreign DNA Should be cut with Same RE

V. imp *

GEL ELECTROPHORESIS

movement

: Separate & Isolate
desired DNA fragment

↓
Electric current

+ → Anode
- → Cathode


* Agarose ^(medium) Gel

(Natural Polymer; Polysaccharide)
(from Gelidium & Gracilaria)

Red Algae
Sea weeds

→ Provide Sieving effect





- * Separation on basis of Size | mol. weight | density
- * Smaller the frag, farther it goes/moves

↓
PYQ Ethidium Bromide, followed
by UV rays

↓
PYQ Bright Orange bands

↓
PYQ ELUTION [Desired frag. cut off from
Agarose gel and Extracted from
gel piece]

9.2.1 Restriction Enzymes

In the year 1963, the two enzymes responsible for restricting the growth of bacteriophage in *Escherichia coli* were isolated. One of these added methyl groups to DNA, while the other cut DNA. The later was called **restriction endonuclease**.

The first restriction endonuclease—*Hind II*, whose functioning depended on a specific DNA nucleotide sequence was isolated and characterised five years later. It was found that *Hind II* always cut DNA molecules at a particular point by recognising a specific sequence of six base pairs. This specific base sequence is known as the **recognition sequence** for *Hind II*. Besides *Hind II*, today we know more than 900 restriction enzymes that have been isolated from over 230 strains of bacteria each of which recognise different recognition sequences.

The convention for naming these enzymes is the first letter of the name comes from the genus and the second two letters come from the species of the prokaryotic cell from which they were isolated, e.g., EcoRI comes from *Escherichia coli* RY 13. In EcoRI, the letter 'R' is derived from the name of

strain. Roman numbers following the names indicate the order in which the enzymes were isolated from that strain of bacteria.

Restriction enzymes belong to a larger class of enzymes called **nucleases**. These are of two kinds; **exonucleases** and **endonucleases**. Exonucleases remove nucleotides from the ends of the DNA whereas, endonucleases make cuts at specific positions within the DNA.

Each restriction endonuclease functions by 'inspecting' the length of a DNA sequence. Once it finds its specific recognition sequence, it will bind to the DNA and cut each of the two strands of the double helix at specific points in their sugar-phosphate backbones (Figure 9.1). Each restriction endonuclease recognises a specific **palindromic nucleotide sequences** in the DNA.





Figure 9.1 Steps in formation of recombinant DNA by action of restriction endonuclease enzyme - EcoRI


Do you know what palindromes are? These are groups of letters that form the same words when read both forward and backward, e.g., “MALAYALAM”. As against a word-palindrome where the same word is read in both directions, the palindrome in DNA is a sequence of base pairs that reads same on the two strands when orientation of reading is kept the same. For example, the following sequences reads the same on the two strands in $5' \rightarrow 3'$ direction. This is also true if read in the $3' \rightarrow 5'$ direction.



Restriction enzymes cut the strand of DNA a little away from the centre of the palindrome sites, but between the same two bases on the opposite strands. This leaves single stranded portions at the ends. There are overhanging stretches called sticky ends on each strand (Figure 9.1). These are named so because they form hydrogen bonds with their complementary cut counterparts. This stickiness of the ends facilitates the action of the enzyme DNA ligase.

Restriction endonucleases are used in genetic engineering to form 'recombinant' molecules of DNA, which are composed of DNA from different sources/genomes.

When cut by the same restriction enzyme, the resultant DNA fragments have the same kind of 'sticky-ends' and, these can be joined together (end-to-end) using DNA ligases (Figure 9.2).



FOR NOTES & DPP CHECK DESCRIPTION

You may have realised that normally, unless one cuts the vector and the source DNA with the same restriction enzyme, the recombinant vector molecule cannot be created.

Separation and isolation of DNA fragments : The cutting of DNA by restriction endonucleases results in the fragments of DNA. These fragments can be separated by a technique known as **gel electrophoresis**. Since DNA fragments are negatively charged molecules they can be separated by forcing them to move towards the anode under an electric field through a medium/matrix. Nowadays the most commonly used matrix is agarose which is a natural polymer extracted from sea weeds. The DNA fragments separate (resolve) according to their size through sieving effect provided by the agarose gel. Hence, the smaller the fragment size, the farther it moves. *Look at the Figure 9.3 and guess at which end of the gel the sample was loaded.*




Figure 9.3 A typical agarose gel electrophoresis showing migration of undigested (lane 1) and digested set of DNA fragments (lane 2 to 4)

The separated DNA fragments can be visualised only after staining the DNA with a compound known as ethidium bromide followed by exposure to UV radiation (you cannot see pure DNA fragments in the visible light and without staining). You can see bright orange coloured bands of DNA in a ethidium bromide stained gel exposed to UV light (Figure 9.3). The separated bands of DNA are cut out from the agarose gel and extracted from the gel piece. This step is known as elution. The DNA fragments purified in this way are used in constructing recombinant DNA by joining them with cloning vectors.



* VECTORS = Gene taxi / Gene vehicle
= CLONING VECTOR

① ori → control Copy no.


2 most commonly used Vectors:-

① Plasmids → 1-2 copy / per cell
→ 15-100 or even higher / cell

② Bacteriophage → high copy no.

② rop : (genes), coding for Proteins, involved in Replication

③ Cloning site
Recognition
RE




If a vector has cloning sites for more RE, it is preferred, as we have more options



* P40 For a specific RE,

Cloning sites should be
very few, preferably simple



↓ Because

> 1 clon. sites generate
multiple fragments


↓
Complicate gene cloning

V.V. Singh



Selectable Marker (genes) :-

- a) Antibiotic resis. gene
- b) Lac-Z gene



• Genes in vector, help in identifying Non-Recombinant and eliminating Transformed from Non-transf.

↓

Selective growth of Transf. (Recomb.)

①

Antibiotic - resist. gene :

Cumbersome process

Ampicillin


Tetracyclin

Chloramphenicol

Kanamycin

E. coli
not
resistant







- a) 8 RE
- b) 2 selec-markers
amp^r, tet^r
- c) ori
- d) rop


pBR322 (1st artificial vector / Plasmid)
 ↓ Boliver & Rodriguez
 plasmid Exp. m.

* Suppose foreign DNA & pBR322 cut with BamHI

E. coli Host / amp^r

- a) amp^r  Transformed Recombinant *
- b) amp^r  Transformed Non-Recomb. *
- c) Non-transf. Non-Recomb.

 ∅DNA formed
amp^r ✓
tet^r X

 amp^r ✓
tet^r ✓
vector
Ampicillin

No ∅DNA formed
Tetracycline

* alive }
* alive }

* die
* alive

die

die



Ques.

If I want Recomb. host to die in
Amp. medium, which RE should be used?


↓
Pst I

or

Pvu I




Ques: pBR322 cut with Sal I;




- a) Recombinant Host → Ampicillin : Alive / die?
- b) Non-transformed Host → Tet : Alive / die?

Ques: If pBR322 cut with Pvu II;



- a) Recombin. Host → Amp : Alive / die?
- b) Non-transf. " → Amp : Alive / die?

⑥ Lac-Z gene S. marker
(Blue-white selection)



Lac-Z-gene

β -galactosidase enzyme

Lactose
(β -galactoside)

Glucose + Galactose
Blue Colonies

<u>E. coli Host</u>		Chr. subs. + Ampicillin
a)	<u>Transformed Recomb</u>	→ <u>alive, white colonies</u>
b)	<u>Transf. Non-Recomb</u>	→ <u>alive, Blue colonies</u>
c)	<u>Non-transformed</u>	→ <u>Died</u>

Chromogenic Substrate

→ due to insertional activation of Lac-Z gene



* Vectors for Plants and Animals :-

↓
Natural Genetic Engineering
* Disarmed
* Modified Ti-plasmid of A. tumefaciens,
no more pathogenic to plant
but still able to deliver
piece of desired DNA

↓
Disarmed/modified/
No more pathogenic
* Retroviruses

* Agrobacterium tumefaciens, a soil bacterium




Tumor inducing
Ti plasmid

T-DNA

Pathogenic to dicot

Crown gall tumor

Ti-plasmid



dicot
plant

9.2.2 Cloning Vectors

You know that plasmids and bacteriophages have the ability to replicate within bacterial cells independent of the control of chromosomal DNA. Bacteriophages because of their high number per cell, have very high copy numbers of their genome within the bacterial cells. Some plasmids may have only one or two copies per cell whereas others may have 15-100 copies per cell. Their numbers can go even higher. If we are able to link an alien piece of DNA with bacteriophage or plasmid DNA, we can multiply its numbers equal to the copy number of the plasmid or bacteriophage. Vectors used at present, are engineered in such a way that they help easy linking of foreign DNA and selection of recombinants from non-recombinants.

The following are the features that are required to facilitate cloning into a vector.

- (i) **Origin of replication (ori)** : This is a sequence from where replication starts and any piece of DNA when linked to this sequence can be made to replicate within the host cells. This sequence is also responsible for controlling the copy number of the linked DNA. So, if one wants to recover many copies of the target DNA it should be cloned in a vector whose origin support high copy number.
- (ii) **Selectable marker** : In addition to 'ori', the vector requires a selectable marker, which helps in identifying and eliminating non-transformants and selectively permitting the growth of the transformants. **Transformation** is a procedure through which a piece of DNA is introduced in a host bacterium (you will study the process in subsequent section). Normally, the genes encoding resistance to antibiotics such as ampicillin, chloramphenicol, tetracycline or kanamycin, etc., are considered useful selectable markers for *E. coli*. The normal *E. coli* cells do not carry resistance against any of these antibiotics.

recomb

- (iii) **Cloning sites:** In order to link the alien DNA, the vector needs to have very few, preferably single, **recognition sites** for the commonly used restriction enzymes. Presence of more than one recognition sites within the vector will generate several fragments, which will complicate the gene cloning (Figure 9.4). The ligation of alien DNA is carried out at a restriction site present in one of the two **antibiotic resistance** genes. For example, you can ligate a foreign DNA at the **BamH I** site of tetracycline resistance gene in the vector pBR322. The recombinant plasmids will lose tetracycline resistance due to insertion of foreign DNA but can still be selected out from non-recombinant ones by plating the transformants on




Figure 9.4 *E. coli* cloning vector pBR322 showing restriction sites (*Hind* III, *Eco*R I, *Bam*H I, *Sal* I, *Pvu* II, *Pst* I, *Cla* I), *ori* and antibiotic resistance genes (*amp*^r and *tet*^r). *rop* codes for the proteins involved in the replication of the plasmid.

tetracycline containing medium. The transformants growing on ampicillin containing medium are then transferred on a medium containing tetracycline. The recombinants will grow in ampicillin containing medium but not on that containing tetracycline. But, non-recombinants will grow on the medium containing both the antibiotics. In this case, one antibiotic resistance gene helps in selecting the transformants, whereas the other antibiotic resistance

amp

tet

gene gets 'inactivated due to insertion' of alien DNA, and helps in selection of recombinants.

Selection of recombinants due to inactivation of antibiotics is a cumbersome procedure because it requires simultaneous plating on two plates having different antibiotics. Therefore, alternative selectable markers have been developed which differentiate recombinants from non-recombinants on the basis of their ability to produce colour in the presence of a chromogenic substrate. In this, a recombinant DNA is inserted within the coding sequence of an enzyme, β -galactosidase. This results into inactivation of the gene for synthesis of this enzyme, which is referred to as **insertional inactivation**. The presence of a chromogenic substrate gives blue coloured colonies if the plasmid in the bacteria does not have an

insert. Presence of insert results into insertional inactivation of the β -galactosidase gene and the colonies do not produce any colour, these are identified as recombinant colonies.


- (iv) **Vectors for cloning genes in plants and animals** : You may be surprised to know that we have learnt the lesson of transferring genes into plants and animals from bacteria and viruses which have known this for ages – how to deliver genes to transform eukaryotic cells and force them to do what the bacteria or viruses want. For example, *Agrobacterium tumefaciens*, a pathogen of several dicot plants is able to deliver a piece of DNA known as 'T-DNA' to transform normal plant cells into a **tumor** and direct these tumor cells to produce the chemicals required by the pathogen. Similarly, retroviruses in animals have the ability to transform normal cells into **cancerous** cells. A better understanding of the art of delivering genes by pathogens in

their eukaryotic hosts has generated knowledge to transform these tools of pathogens into useful vectors for delivering genes of interest to humans. The tumor inducing (Ti) plasmid of *Agrobacterium tumifaciens* has now been modified into a cloning vector which is no more pathogenic to the plants but is still able to use the mechanisms to deliver genes of our interest into a variety of plants. Similarly, retroviruses have also been disarmed and are now used to deliver desirable genes into animal cells. So, once a gene or a DNA fragment has been ligated into a suitable vector it is transferred into a bacterial, plant or animal host (where it multiplies).

9.2.3 Competent Host (For Transformation with Recombinant DNA)

① Since DNA is a hydrophilic molecule, it cannot pass through cell membranes. Why? In order to force bacteria to take up the plasmid, the bacterial cells must first be made 'competent' to take up DNA. This is done by treating them with a specific concentration of a divalent cation, such as calcium, which increases the efficiency with which DNA enters

hydrophobic tails



the bacterium through pores in its cell wall. Recombinant DNA can then be forced into such cells by incubating the cells with recombinant DNA on ice, followed by placing them briefly at 42°C (heat shock), and then putting them back on ice. This enables the bacteria to take up the recombinant DNA.

This is not the only way to introduce alien DNA into host cells. In a method known as **micro-injection**, recombinant DNA is directly injected into the nucleus of an animal cell. In another method, suitable for plants, cells are bombarded with high velocity micro-particles of gold or tungsten coated with DNA in a method known as **biolistics** or **gene gun**. And the last method uses 'disarmed pathogen' vectors, which when allowed to infect the cell, transfer the recombinant DNA into the host.

Now that we have learnt about the tools for constructing recombinant DNA, let us discuss the processes facilitating recombinant DNA technology.

- ⑤ Polyethylene Glycol (PEG)
Chemofusion
- ⑥ Electrofusion



240

* Processes of Biotechnology :-

1. Isolation of Genetic Material (DNA) :

Lyases, chilled Ethanol, Spooling

2. Cutting of DNA at Specific Locations :

by Restriction Endonuclease

3. Separation and Isolation of Desired DNA fragment :

by Gel Electrophoresis
[Ethidium Br → UV rays → Bright Orange Bands]
ELUTION

4. Amplification of Gene of Interest : by PCR

5. λDNA Construction : by DNA Ligase

6. Insertion of λDNA into Competent Host

Heat-shock, Gene Gun, Microneedle, Chemical Electroporation

Disarmed Vectors like Retrovirus
Ti-plasmid of A. tumefaciens

Use of Selectable Marker [Identify Transformed, Recombinant] HOST



- 7. Obtaining the Foreign Gene Product
- 8. Downstream processing



* PCR = Polymerase Chain Reaction (developed by Kary Mullis)

PYQs


• Amplification of GOI
Multiple identical copies of GOI, in-vitro

* Denaturation → Annealing → Elongation
Tag polymerase

* Thermal Cycles



After 30 cycles,
≈ 1 billion copies



I Denaturation ↓ 90-95°C Heat
H-bonds break and both strands separate

II Annealing + 3' Primer addition at 3' end
50-65°C

III EXTENSION
ELONGATION
Polymerisation
72°C

↓ dNTPs (activated nucleotides)
↓ Taq polymerase (DNA Polymerase)

- Thermus aquaticus bacteria
- Thermostable enzyme
- Active at high temp. induced denaturation of dsDNA

9.3 PROCESSES OF RECOMBINANT DNA TECHNOLOGY

Recombinant DNA technology involves several steps in specific sequence such as isolation of DNA, fragmentation of DNA by restriction endonucleases, isolation of a desired DNA fragment, ligation of the DNA fragment into a vector, transferring the recombinant DNA into the host, culturing the host cells in a medium at large scale and extraction of the desired product. Let us examine each of these steps in some details.



————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

9.3.1 Isolation of the Genetic Material (DNA)

Recall that nucleic acid is the genetic material of all organisms without exception. In majority of organisms this is deoxyribonucleic acid or DNA. In order to cut the DNA with restriction enzymes, it needs to be in pure form, free from other macro-molecules. Since the DNA is enclosed within the membranes, we have to break the cell open to release DNA along with other macromolecules such as RNA, proteins, polysaccharides and also lipids. This can be achieved by treating the bacterial cells/plant or animal tissue with enzymes such as **lysozyme** (bacteria), **cellulase** (plant cells), **chitinase** (fungus). You know that genes are located on long molecules of DNA



Figure 9.5 DNA that separates out can be removed by spooling

intertwined with proteins such as histones. The RNA can be removed by treatment with ribonuclease whereas proteins can be removed by treatment with protease. Other molecules can be removed by appropriate treatments and purified DNA ultimately precipitates out after the addition of chilled ethanol. This can be seen as collection of fine threads in the suspension (Figure 9.5).

9.3.2 Cutting of DNA at Specific Locations

Restriction enzyme digestions are performed by incubating purified DNA molecules with the restriction enzyme, at the optimal conditions for that specific enzyme. Agarose gel electrophoresis is employed to check the progression of a restriction enzyme digestion. DNA is a negatively charged molecule, hence it moves towards the positive electrode (anode) (Figure 9.3). The process is repeated with the vector DNA also.

The joining of DNA involves several processes. After having cut the source DNA as well as the vector DNA with a specific restriction enzyme, the cut out 'gene of interest' from the source DNA and the cut vector with space are mixed and ligase is added. This results in the preparation of recombinant DNA.

9.3.3 Amplification of Gene of Interest using PCR

PCR stands for **Polymerase Chain Reaction**. In this reaction, multiple copies of the gene (or DNA) of interest is synthesised *in vitro* using two




Figure 9.6 Polymerase chain reaction (PCR) : Each cycle has three steps: (i) Denaturation; (ii) Primer annealing; and (iii) Extension of primers

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

sets of primers (small chemically synthesised oligonucleotides that are complementary to the regions of DNA) and the enzyme DNA polymerase. The enzyme extends the primers using the nucleotides provided in the reaction and the genomic DNA as template. If the process of replication of DNA is repeated many times, the segment of DNA can be amplified to approximately billion times, i.e., 1 billion copies are made. Such repeated amplification is achieved by the use of a thermostable DNA polymerase (isolated from a bacterium, *Thermus aquaticus*), which remain active during the high temperature induced denaturation of double stranded DNA. The amplified fragment if desired can now be used to ligate with a vector for further cloning (Figure 9.6).

9.3.4 Insertion of Recombinant DNA into the Host Cell/Organism

There are several methods of introducing the ligated DNA into recipient cells. Recipient cells after making them 'competent' to receive, take up DNA present in its surrounding. So, if a recombinant DNA bearing gene for resistance to an antibiotic (e.g., ampicillin) is transferred into *E. coli* cells, the host cells become transformed into ampicillin-resistant cells. If we spread the transformed cells on agar plates containing ampicillin, only transformants will grow, untransformed recipient cells will die. Since, due to ampicillin resistance gene, one is able to select a transformed cell in the presence of ampicillin. The ampicillin resistance gene in this case is called a **selectable marker**.



9.3.5 Obtaining the Foreign Gene Product

When you insert a piece of alien DNA into a cloning vector and transfer it into a bacterial, plant or animal cell, the alien DNA gets multiplied. In almost all recombinant technologies, the ultimate aim is to produce a desirable protein. Hence, there is a need for the recombinant DNA to be expressed. The foreign gene gets expressed under appropriate conditions. The expression of foreign genes in host cells involve understanding many technical details.

After having cloned the gene of interest and having optimised the conditions to induce the expression of the target protein, one has to

consider producing it on a large scale. Can you think of any reason why there is a need for large-scale production? If any protein encoding gene is expressed in a heterologous host, it is called a **recombinant protein**. The cells harbouring cloned genes of interest may be grown on a small scale in the laboratory. The cultures may be used for extracting the desired protein and then purifying it by using different separation techniques.

The cells can also be multiplied in a continuous culture system wherein the used medium is drained out from one side while fresh medium is added from the other to maintain the cells in their physiologically most

active log/exponential phase. This type of culturing method produces a larger biomass leading to higher yields of desired protein.

Small volume cultures cannot yield appreciable quantities of products. To produce in large quantities, the development of **bioreactors**, where large volumes (100-1000 litres) of culture can be processed, was required. Thus, bioreactors can be thought of as vessels in which raw materials are biologically converted into specific products, individual enzymes, etc., using microbial plant, animal or human cells. A bioreactor provides the optimal conditions for achieving the desired product by providing optimum growth conditions (temperature, pH, substrate, salts, vitamins, oxygen).

The most commonly used bioreactors are of stirring type, which are shown in Figure 9.7.




Figure 9.7 (a) Simple stirred-tank bioreactor; (b) Sparged stirred-tank bioreactor through which sterile air bubbles are sparged

A stirred-tank reactor is usually cylindrical or with a curved base to facilitate the mixing of the reactor contents. The stirrer facilitates even mixing and oxygen availability throughout the bioreactor. Alternatively air can be bubbled through the reactor. If you look at the figure closely you will see that the bioreactor has an agitator system, an oxygen delivery system and a foam control system, a temperature control system, pH control system and sampling ports so that small volumes of the culture can be withdrawn periodically.


9.3.6 Downstream Processing

After completion of the biosynthetic stage, the product has to be subjected through a series of processes before it is ready for marketing as a finished

product. The processes include separation and purification, which are collectively referred to as downstream processing. The product has to be formulated with suitable preservatives. Such formulation has to undergo thorough clinical trials as in case of drugs. Strict quality control testing for each product is also required. The downstream processing and quality control testing vary from product to product.


Culture — 2 types

I ^{NCERT} CONTINUOUS CULTURE SYSTEM [Open System]



- ^{NCERT} { • Used medium is drained from one end while fresh medium is added continuously from other end

II BATCH Culture System [Closed System]




- No new fresh medium can be added in between and used medium drained only at end of process.

NCERT

• Maintain Cells in
(Host)
their physiologically
Most Active phase


i.e. * log / Exponential phase



↓
Produces larger Biomass

↓
↑↑ Yield of Desired
Protein

- Lesser yield of desired product
- 3 phases - lag / log / Stable.



QUESTION – 01 (2015 Re)

The cutting of DNA at specific locations became possible with the discovery of:

- A** Probes
- B** Selectable markers
- C** Ligases
- D** Restriction enzymes

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

QUESTION – 02 (2022)

In the following palindromic base sequences of DNA, which one can be cut easily by particular restriction enzyme?

- A** 5' GTATTC 3' ; 3' CAT AAG 5'
- B** 5' GATACT 3' ; 3' CTATGA 5'
- C** 5' GAATTC 3' ; 3' CTTAAG 5'
- D** 5' CTCAGT 3' ; 3' GAGTCA 5'

————— FOR NOTES & DPP CHECK DESCRIPTION —————

QUESTION – 03 (2022)

Which of the following is not a desirable feature of a cloning vector?

false

- A** Presence of two or more recognition sites
- B** Presence of origin of replication
- C** Presence of a marker gene
- D** Presence of single restriction enzyme site

————— FOR NOTES & DPP CHECK DESCRIPTION —————

QUESTION – 04 (2022)



Given below are two statements:

Statement I: Restriction endonucleases recognise specific sequence to cut DNA known as palindromic nucleotide sequence.

Statement II: Restriction endonucleases cut the DNA strand a little away from the centre of the palindromic site.

In the light of the above statements, choose the most appropriate answer from the options given below.

A Statement I is incorrect but Statement II is correct

B Both Statement I and Statement II are correct

C Both statement I and statement II are incorrect

D Statement I is correct but Statement II is incorrect

FOR NOTES & DPP CHECK DESCRIPTION

QUESTION – 05 (2021)

DNA strands on a gel stained with ethidium bromide when viewed under UV radiation, appear as:

- ☒ **A** Bright orange bands
- ☐ **B** Dark red bands
- ☐ **C** Bright blue bands
- ☐ **D** Yellow bands

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

QUESTION – 06 (2021)



Plasmid pBR322 has PstI restriction enzyme site within gene amp^R that confers ampicillin resistance, If this enzyme is used for inserting a gene for β -galactoside production and the recombinant plasmid is inserted in an E.coli strain. (GDI)

- A** The transformed cells will have the ability to resist ~~ampicillin~~ as well as produce β -galactoside.
 Recomb. *amp^R X*
- B** It will lead to lysis of host cell.
 X
- C** It will be able to produce a novel protein with dual ability.
 X
- D** It will not be able to confer ampicillin resistance to host cell.
 X

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 07 (2021)

A specific recognition sequence identified by endonucleases to make cuts at specific positions within the DNA is:

- A** Okazaki sequences ~~X~~
- B** ~~✓~~ Palindromic Nucleotide sequences
- C** Poly (A) tail sequences ~~X~~
- D** Degenerate primer sequence ~~X~~

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

QUESTION – 08 (2020-Covid)

First discovered restriction endonuclease that always cuts DNA molecule at a particular point by recognising a specific sequence of six base pairs is:

- A** Adenosine deaminase
- B** Thermostable DNA polymerase
- C** Hind II
- D** EcoRI

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 09 (2020)

Match the organism with its use in biotechnology.
Select the correct option from the following:

- A** A-(IV), B-(III), C-(I), D-(II)
- B** A-(III), B-(II), C-(IV), D-(I)
- C** A-(III), B-(IV), C-(I), D-(II)
- D** A-(II), B-(IV), C-(III), D-(I)

(A)	<i>Bacillus thuringiensis</i>	(I)	Cloning vector
(B)	<i>Thermus aquaticus</i>	(II)	Construction of first rDNA molecule
(C)	<i>Agrobacterium tumefaciens</i>	(III)	DNA polymerase
(D)	<i>Salmonella typhimurium</i>	(IV)	Cry proteins

QUESTION – 10 (2020)

The sequence that controls the copy number of the linked DNA in the vector, is termed:

- A** ~~Ori site~~
- B** Palindromic sequence
- C** Recognition site
- D** Selectable marker

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 11 (2020)

The specific palindromic sequence which is recognized by EcoRI is:

- A** 5' – GGAACC – 3' ; 3' – CCTTGG – 5'
- B** 5' – CTTAAG – 3' ; 3' – GAATTC – 5'
- C** 5' – GGATCC – 3' ; 3' – CCTAGG – 5'
- D** ~~5' – GAATTC – 3' ; 3' – CTTAAG – 5'~~

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 12 (2020)

Identify the wrong statement with regard to restriction enzymes.

- A** They cut the strand of DNA at palindromic sites. *false*
- B** They are useful in genetic engineering. ✓
- C** ~~Sticky ends can be joined by using DNA ligases.~~ *True for*
- D** Each restriction enzyme functions by inspecting the length of a DNA sequence. ✓

————— FOR NOTES & DPP CHECK DESCRIPTION —————

QUESTION – 13 (2020)

Choose the correct pair from the following:

- A** Polymerases – Break the DNA into fragments *RE*
- B** Nucleases – Separate the two strands of DNA *Heli*
- C** Exonucleases – Make cuts at specific positions within DNA
- D** Ligases – Join the two DNA molecules

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

QUESTION – 14 (2019)

Following statements describe the characteristics of the enzyme restriction endonuclease. Identify the incorrect statement. *false*

- A** The enzyme cuts DNA molecule at identified position within the DNA. ✓
- B** The enzyme binds DNA at specific sites and cuts only one of the two strands. ~~both~~ ✗
- C** The enzyme cuts the sugar-phosphate backbone at specific sites on each strand. ✓
- D** The enzyme recognizes a specific palindromic nucleotide sequence in the DNA. ✓

QUESTION – 15 (2017-Delhi)

A gene whose expression helps to identify transformed cell is known as

- A** Selectable marker
- B** Vector
- C** Plasmid
- D** Structural gene

————— FOR NOTES & DPP CHECK DESCRIPTION —————

QUESTION – 16 (2017)

Restriction endonucleases are:

- A** Used in genetic engineering for ligating two DNA molecules *DNA ligase*
- B** Used for in vitro DNA synthesis *PCR*
- C** Synthesised by bacteria as part of their defense mechanism
- D** Present in mammalian cell for degradation of DNA when the cell dies

————— FOR NOTES & DPP CHECK DESCRIPTION —————

QUESTION – 17 (OS, 2016-II)



Add in notes


Which of the following restriction enzymes produces blunt ends?

- A** Xho I
- B** Hind III
- C** Sal I
- D** Eco RV

OUT OF NCERT



Sma I



Blunt ends

FOR NOTES & DPP CHECK DESCRIPTION

QUESTION – 18 (2016-II)

A foreign DNA and plasmid cut by the same restriction endonuclease can be joined to form a recombinant plasmid using:

- A** Polymerase-III
- B** Ligase
- C** Eco RI
- D** Taq polymerase

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 19 (2016-I)

Which of the following is a restriction endonuclease?

- A** ~~Hind II~~
- B** Protease *lyases*
- C** DNase I
- D** RNase *lyases*

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 20 (2016-I)

Which of the following is not a feature of the plasmids?

- A** Independent replication
- B** Circular structure
- C** Transferable
- D** Single-stranded

false

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 21 (2015 Re)

The DNA molecule to which the gene of interest is integrated for cloning is called:

- A** Vector
- B** Template
- C** Carrier
- D** Transformer

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 22 (2015 Re)

The introduction of t-DNA into plants involves:

- A** Altering the pH of the soil, then heat shocking the plants
- B** Exposing the plants to cold for a brief period
- C** Allowing the plant roots to stand in water
- D** Infection of the plant by *Agrobacterium tumefaciens*

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 23 (2014)

Which vector can clone only a small fragment of DNA?

- A** Cosmid ~~X~~
- B** Bacterial artificial chromosome
- C** Yeast artificial chromosome
- D** Plasmid

HGP
in Mol. basis

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 24 (2013)

DNA fragments generated by the restriction endonuclease in a chemical reaction can be separated by:

- A** Restriction mapping
- B** Centrifugation
- C** Polymerase chain reaction
- D** Electrophoresis

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 25 (2013)



The colonies of recombinant bacteria appear white in contrast to blue colonies of non-recombinant bacteria because of:

lacZ gene inactive → β -galac

Insert inact of β -galac. gene

X galactosidase

(may be non-transf.)

X

- A** Inactivation of glycosidase enzyme in recombinant bacteria
- B** Non-recombinant bacteria containing betagalactosidase
- C** Insertional inactivation of alpha-galactosidase in non-recombinant bacteria
- D** Insertional inactivation of beta-galactosidase in recombinant bacteria ✓

FOR NOTES & DPP CHECK DESCRIPTION

QUESTION – 26 (2022)

Which one of the following statement is **not true** regarding **gel electrophoresis** technique?

- A** Bright orange coloured bands of DNA can be observed in the gel when exposed to UV light. ✓
- B** The process of extraction of separated DNA strands from gel is called elution. ✓
- C** The separated DNA fragments are stained by using ethidium bromide. ✓
- D** ~~The presence of chromogenic substrate gives blue coloured DNA bands on the gel~~ ✗

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

QUESTION – 27 (2022)

Given below are two statements: one is labelled as Assertion (A) and the other is labelled as Reason (R).

Assertion (A): Polymerase chain reaction is used in DNA amplification. ✓ ^{PCR}

Reason (R): The ampicillin resistant gene is used as a selectable marker to check transformation. ✓ _{vector}

In the light of the above statements, choose the correct answer from the options given below.

- ☒ **A** (A) is not correct but (R) is correct
- ☒ **B** Both (A) and (R) are correct and (R) is the correct explanation of (A)
- ☒ **C** Both (A) and (R) are correct but (R) is not the correct explanation of (A)
- ☒ **D** (A) is correct but (R) is not correct

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

QUESTION – 28 (2021)

During the purification process for recombinant DNA technology, addition of chilled ethanol precipitates out:

- ☒ **A** DNA
- ☐ **B** Histones
- ☐ **C** Polysaccharides
- ☐ **D** RNA

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 29 (2021)

During the process of gene amplification using PCR, if very high temperature is not maintained in the beginning, then which of the following steps of PCR will be affected first?

- A** Extension
- B** Denaturation
- C** Ligation
- D** Annealing

Denat
90-95°C

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 30 (2020)





In gel electrophoresis, separated DNA fragments can be visualized with the help of:

- A** Ethidium bromide in UV radiation
- B** Acetorarmine in UV radiation
- C** Ethidium bromide in infrared radiation
- D** Acetocarmine in bright blue light

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 31 (2020-Covid)

In a mixture, DNA fragments are separated by

- A** Restriction digestion 
- B** Electrophoresis 
- C** Polymerase chain reaction 
- D** Bioprocess engineering 

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 32 (2020-Covid)

In recombinant DNA technology antibiotics are used:

- A** To detect alien DNA
- B** To impart disease-resistance to the host plant
- C** As selectable markers
- D** To keep medium bacteria-free

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 33 (2020-Covid)

Match the following techniques or instruments with their usage:
Select the correct option from following:

A ~~A-(II), B-(I), C-(IV), D-(III)~~

B ~~A-(IV), B-(III), C-(II), D-(I)~~

C ~~A-(II), B-(I), C-(III), D-(IV)~~

D ~~A-(III), B-(II), C-(IV), D-(I)~~

(A)	Bioreactor	(I)	Separation of DNA fragments
(B)	Electrophoresis	(II)	Production of large quantities of products
(C)	PCR	(III)	Detection of pathogen, based on antigen-antibody reaction
(D)	ELISA	(IV)	Amplification of nucleic acids

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

QUESTION – 34 (2020-Covid)

Spooling is:

- A** Cutting of separated DNA bands from the agarose gel → Elution
- B** Transfer of separated DNA fragments to synthetic membranes X Southern Blotting
- C** Collection of isolated DNA ↗
- D** Amplification of DNA PCR

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 35 (2020-Covid)

Select the correct statement from the following:

- A** The polymerase ~~enzyme~~ ^{ligase} joins the gene of interest and the vector DNA
- B** ~~Restriction enzyme~~ digestions are performed by incubating purified DNA molecules with the restriction enzymes of optimum conditions
- C** ~~PCR~~ ^{Gel Electroph} is used for isolation and separation of gene of interest
- D** Gel electrophoresis ~~is~~ ^{PCR} used for amplification of a DNA segment

————— FOR NOTES & DPP CHECK DESCRIPTION —————

QUESTION – 36 (2019)

Which one of the following equipments is essentially required for growing microbes on a large scale, for industrial production of enzymes?

- A** BOD incubator
- B** Sludge digester
- C** Industrial oven
- D** Bioreactor

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 37 (2019)

DNA precipitation out of a mixture of biomolecules can be achieved by treatment with

- A** Isopropanol
- B** ~~Chilled ethanol~~
- C** Methanol at room temperature
- D** Chilled chloroform

———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

QUESTION – 38 (2018)

The correct order of steps in Polymerase Chain Reaction (PCR) is:

- A** Extension, Denaturation, Annealing
- B** Annealing, Extension, Denaturation
- C** Denaturation, Extension, Annealing
- D** Denaturation, Annealing, Extension

———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

QUESTION – 39 (2017-Delhi)

What is the criterion for DNA fragments movement on agarose gel during gel electrophoresis?

- A** The larger the fragment size, the farther it moves
- B** ~~The smaller the fragment size, the farther it moves~~
- C** ~~Positively charged fragments move to farther end~~
- D** ~~Negatively charged fragments do not move~~

———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

QUESTION – 40 (2017-Delhi)

The process of separation and purification of expressed protein before marketing is called

- A** Upstream processing
- B** Downstream processing
- C** Bioprocessing
- D** Post-production processing

———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

QUESTION – 41 (2017-Delhi)

The DNA fragments separated on an agarose gel can be visualised after staining with:

- A** Bromophenol blue
- B** Acetocarmine
- C** Aniline blue
- D** Ethidium bromide

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 42 (2016-II)

Which of the following is not a component of downstream processing?

- A** Preservation ✓
- B** Expression ✗
- C** Separation ✓
- D** Purification ✓

———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

QUESTION – 43 (2016-II)

Stirred-tank bioreactors have been designed for:

- A** Availability of oxygen throughout the process
- B** Ensuring anaerobic conditions in the culture vessel
- C** Purification of product
- D** Addition of preservatives to the product

Downstream
Downst

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 44 (2016-I)

The Taq polymerase enzyme is obtained from:

- A** *Thermus aquaticus*
- B** *Thiobacillus ferrooxidans*
- C** *Bacillus subtilis*
- D** *Pseudomonas putida*

————— FOR NOTES & DPP CHECK DESCRIPTION —————

QUESTION – 45 (2014)

In vitro clonal propagation in plants is characterised by:

- ☐ A Microscopy
- ☒ B PCR and RAPD
- ☐ C Northern blotting
- ☐ D Electrophoresis and HPLC

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 46 (2013)

Which of the following is not correctly matched for the organism and its cell wall degrading enzyme?

- A** Fungi – Chitinase ✓
- B** Bacteria – Lysozyme ✓
- C** Plant cells – Cellulase ✓
- D** ~~Algae – Methylase~~ ✗

false

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 47 (2023)

Upon exposure to UV radiation, DNA stained with ethidium bromide will show

- A** Bright orange colour
- B** Bright red colour
- C** Bright blue colour
- D** Bright yellow colour

———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

QUESTION – 48 (2023)

In gene gun method used to introduce alien DNA into host cells, microparticles of _____ metal are used.

- A** Silver
- B** Copper
- C** Zinc
- D** Tungsten or gold

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 49 (2023)

During the purification process for recombinant DNA technology, addition of chilled ethanol precipitates out

- A** Polysaccharides
- B** RNA
- C** DNA
- D** Histones

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 50 (2023)

Main steps in the formation of Recombinant DNA are given below. Arrange these steps in a correct sequence

- A. Insertion of recombinant DNA into the host cell. (4)
- B. Cutting of DNA at specific location by restriction enzyme. (1)
- C. Isolation of desired DNA fragment. (2)
- D. Amplification of gene of interest using PCR. (3)

Choose the correct answer from the option given below:

A ~~B, D, A, C~~

B ~~B, C, D, A~~

C C, A, B, D

D C, B, D, A

FOR NOTES & DPP CHECK DESCRIPTION

QUESTION – 51 (2023)

Which of the following is not a cloning vector?

- A** Probe ☒
- B** BAC ☒
- C** YAC ☒
- D** pBR322 ☒

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 52 (NEET-2024)

Hind II always cuts DNA molecules at a particular point called recognition sequence and it consists of:

- A** 4 bp
- B** 10 bp
- C** 8 bp
- D** 6 bp

———— FOR NOTES & DPP CHECK DESCRIPTION ————

QUESTION – 53 (NEET-2024)



What is the fate of a piece of DNA carrying gene of interest which is transferred into and organism?

- A. The piece of DNA would be able to multiply itself independently in the progeny cells of the organism. ~~X~~
- B. It may get integrated into the genome of the recipient. ✓
- C. It may multiply and be inherited along with the host DNA. ✓
- D. The alien piece of DNA is not an integral part of chromosome. ~~X~~
- E. It shows ability to replicate. ~~X~~

Choose the correct answer from the options given below:

A B and C only ✓

B ~~A and E only~~

C ~~A and B only~~


D ~~D and E only~~

———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

QUESTION – 54 (NEET-2024)

The following diagram showing restriction sites in E. coli cloning vector pBR322. Find the role of 'X' and 'Y' genes:

- A** The gene 'X' is for protein involved in replication of Plasmid and 'Y' for resistance to antibiotics.
- B** Gene 'X' is responsible for recognition sites and 'Y' is responsible for antibiotic resistance.
- C** The gene 'X' responsible for resistance to antibiotics and 'Y' for protein involved in the replication of Plasmid.
- D** The gene 'X' is responsible for controlling the copy number of the linked DNA and 'Y' for protein involved in the replication of plasmid.



————— **FOR NOTES & DPP CHECK DESCRIPTION** —————



Homework



read NCERT
word by word

————— **FOR NOTES & DPP CHECK DESCRIPTION** —————

THANK YOU

