



PRACHAND NEET



ONE SHOT



Organic Chemistry

Hydrocarbon

By – Sudhanshu Sir



Topics

to be covered



1 Alkanes

2 Alkenes

3 Alkynes

4 Benzene



PRACHAND SERIES

TELEGRAM CHANNEL



@PW_YAKEENDROPPER

Basics of Organic chemistry

- ① Nomenclature
- ② Isomerism
- ③ G.O.C

⑮ Practical organic chemistry

Reactions


- ① Alkane
 - ② Alkene
 - ③ Alkyne
 - ④ Benzene
- } Hydrocarbon

- ⑤ $R-X$
 - ⑥ 
- } Haloalkanes & Haloarenes

- ⑦ $R-OH$
 - ⑧ 
 - ⑨ $R-O-R$
- } alcohol, phenol & ether
- ⑩ $R-\overset{\overset{O}{\parallel}}{C}-H$
 - ⑪ $R-\overset{\overset{O}{\parallel}}{C}-R$
 - ⑫ $R-\overset{\overset{O}{\parallel}}{C}-OH$
- } Aldehyde, ketone & Carboxylic acid.
- ⑬ Amines → Amines
 - ⑭ Biomolecules



Introduction




Types of Reactions

① Addition Reaction

Alkanes

unsaturated compounds (D.O.U > 0)




→ D.O.U decreases

→ weak bond dissociates

→ strong bond formation

→ entropy decrease

② Substitution Reaction




Substitution → ESR
NSR

Addition → EAR
NAR

elimination → E¹
E²
E¹CB

③ Elimination Rxn



④ Rearrangement Reaction






Another Classification

Elementary/Single step Reaction

- rxn occurs in one step.
- no mechanism needed.



$$\Delta H = H_P - H_R < 0$$


(Exothermic)

$$\Delta H = H_P - H_R > 0$$

(endothermic)

Complex/Multi-step Reaction

- rxn occurs in more than one step.
- mechanism needed.
- slowest step / High E_a step = Rate determining step.



For 'n' step

- Transition state = n
- Intermediate = (n-1)

Intermediate

⇒ for 'n' step rxn ⇒ Intermediate = (n-1)

⇒ less energy

⇒ more stability

⇒ separable from reaction mixture
or

Rearrangement can be possible

⇒ obeys rules of chemical bonding

⇒ formed after first step and consumed till last step.

Transition State

⇒ for 'n' step rxn, T.S = (n)

⇒ more energy


⇒ less stability

⇒ Cannot be separated from reaction mixture
or

Cannot undergo rearrangement

⇒ do not obey rules of chemical bonding

⇒ formed in each step and consumed.



Intermediates



① Carbocation



→ charged
→ Diamagnetic

② Carbanion



③ Carbon free radical



→ odd e^-
→ paramagnetic
→ neutral

④ Benzynes



⑤ Nitrene



⑥ Carbene



neutral

NOTE

→ out of 6 intermediates, only one intermediate C^+ undergoes rearrangement for gain of stability.

Carbocation



Stability of C^+
↳ Dancing Resonance > Aromaticity > α -effect > +M > +H > +I
 $\Delta-CH_2^+$
antiaromaticity < -M < -H < -I

→ Hybridisation = sp^2

→ 3 σ bond

→ Zero lone pair

→ e^- deficient species


→ electrophile → when attack

→ Diamagnetic

→ vacant 2p orbital

→ trigonal planar

→ stereochemistry → if it has chiral centre, then it will give equal molar of (d) & (l)



Rearrangement of Carbocation

→ Rearrangement is possible when stability increases.



stability remains same }
 stability decreases } No rearrangement of C^+

C^+ Rearrangement

- ① Hydride shift
- ② phenyl shift
- ③ Alkyl shift
- ④ Ring contraction
- ⑤ Ring expansion






Hydride Shift






stability increases.
Rearrangement possible




Mechanism






Bond length decrease
Bond energy increase.



Questions of Hydride Shift




QUESTION





Phenyl Shift




⇒ phenyl shift > alkyl shift

① π -bond breaking up.



② Resonance stabilised.

③ No partial double in T.S.







Alkyl Shift






Rearrangement of C^+

- ① $\text{H} > \text{D} > \text{T}$
- ② $\text{ph-EDG} > \text{ph} > \text{ph-EWG}$
- ③ $\text{3}^\circ > \text{2}^\circ > \text{1}^\circ$ alkyl shift

Questions on Rearrangement:








(4) Ring Expansion



4 membered ring $\rightarrow$ (5)
5 membered ring $\rightarrow$ (6)

Condition $\Rightarrow$ (+ve) charge must be outside the ring.








Dancing Resonance

(5) Ring Contraction








→ 4 membered ring → ③
→ 7 membered ring → ⑥
Condition ⇒ charge ⊕ve on the ring.





H-shift






QUESTION



QUESTION




Hydride
shift →




Free Radical



Formation (Homolytic Fission)



sunlight or $h\nu$
 $T > 500^\circ\text{C}$ or 773K
peroxide catalyst



- 3 σ bonds, Neutral
- Zero lone pair
- Odd e^- molecule

- $e^- \rightarrow$ p-orbital
- Highly unstable
- Highly reactive
- sp^2 hybridized $\rightarrow$ Trigonal Planar
- Paramagnetic, $\mu \neq 0$
- $E^\oplus$, e^- deficient, $7 e^-$

Natural Tendencies of F.R.

free radical $\rightarrow$ odd e^-

highly unstable.



unpaired e^-

$\rightarrow$ high tendency to pair up


Combination





Disproportionation



1. Combination:



2. Disproportionation:



Note:

Degree


1° F.R → always combination.

2° F.R → can show (C) + (D)


3° F.R → only shows (D)

if there is no option of disproportionation,
consider combination products of
all F.R


Temperature





Kinetics $\Rightarrow$ Intermolecular $<$ Intramolecular



Stability $\Rightarrow$ (Resonance stability) $\uparrow$ more preferred.



Hydrocarbon



Introduction

<u>Alkane</u> →	$C_n H_{2n+2}$	$D.O.U = 0$	Saturated Compounds.
Alkene →	$C_n H_{2n}$	$D.O.U = 1$	Unsaturated compounds.
Alkyne →	$C_n H_{2n-2}$	$D.O.U = 2$	" "
Alkadiene/ Dialkene →	$C_n H_{2n-2}$	$D.O.U = 2$	" "
Benzene →	$C_6 H_6$	$D.O.U = 4$	" "

Alkanes



↳ known as Paraffins
(little reactivity)

↳ sp^3 hybridised.


↳ Cannot show addition rxn

↳ shows Free radical substitution.

① Sunlight / $h\nu$

② $T > 500^\circ C$

③ Peroxide Catalyst




Preparations of Alkanes

(a) From Alkyl Halide ($R-X$)

Ar = Aromatic

1. Wurtz Reaction

R = aliphatic



Mechanism → Free radical mechanism

Reactivity order → $R-I > R-Br > R-Cl > R-F$

dry ether is used → ether → non-polar solvent




dry condition → to avoid moisture

Intermediate = $R^\bullet$







→ product = symmetrical alkane

→ methane (CH_4) X





⇒ not preferred for unsymmetrical alkane as it gives mixture of products.




Mechanism:






QUESTION




QUESTION








2. Fitting Reaction




Biphenyl



3. Wurtz fitting Reaction




4. Frankland Reaction



(Frankland Reagent = R_2Zn)




5. Ullmann's Reaction



6. Corey-house Synthesis

~~methane~~


↳ Used for preparation of unsymmetrical alkanes. (R-R')





Preparation of Gilman's Reagent



7. Using Grignard Reagent



methane
preparation
✓✓




Preparation of Grignard reagent



$\Rightarrow$ Grignard reagent is kept anhydrous or prepared in lab at the time of rxn.

$\downarrow$
very reactive to form alkane by taking acidic H^+ .


8. Reduction



9. Red P/HI

↳ Best Reducing agent
↳ BAAP Reagent






- $\text{R-X} \rightarrow$
- ① wurtz
 - ② Fittig
 - ③ wurtz-fittig
 - ④ Frankland
 - ⑤ Ullmann
 - ⑥ Grignard reagents
 - ⑦ Corey house synthesis
 - ⑧ Reduction
 - ⑨ Red P + HI

(b) From Carboxylic Acids



1. Kolbe's Electrolysis





→ free radical mechanism
→ methane X

→ anode → $R-R + CO_2(g)$

→ cathode → $H_2\uparrow + NaOH$



→ pH increase
→ symmetrical alkane





Reaction Mechanism






At anode







At Cathode




Kolbe's Electrolysis:






2. Decarboxylation: (Removal of CO₂)

↳ prefix meaning for Removal



Mechanism



⇒ Rate of decarboxylation $\propto$ stability of Carboanion



→ Intermediate = Carboanion
 R^-

↳ no Rearrangement

↳ Stability order of C^-

Aromaticity > Back Bonding

✓
-M
✓
-H
✓
-I

antiaromaticity

< +M < +H < +I <

QUESTION



Which of the following is correct order for the ease of decarboxylation of these carboxylic acids?

CH_3COOH , CF_3COOH , CCl_3COOH

(a)

(b)


(c)

A $a > b > c$


B $c > b > a$

C $b > c > a$

D $b > a > c$



3. Red P + HI




(C) From Aldehyde & Ketone




1. Red R + HI




2. Wolff Kishner Reduction




→ Alert ⇒ avoid
Base sensitive
group.





Mechanism




3. Clemmensen's Reduction





Alert = avoid Acid sensitive group.



(d) From Aluminium & Beryllium Carbide



(e) From Alkene & Alkyne (Hydrogenation)








$\Delta H_{\text{hydrogenation}} \propto \text{no. of } \pi\text{-bonds} \propto \frac{1}{\text{stability}}$
Always exothermic

- ① $\text{R-X} \rightarrow ①$
- ② $\text{R}-\overset{\text{O}}{\underset{\text{H}}{\text{C}}}-\text{OH} \rightarrow ③$
- ③ $\text{R}-\overset{\text{O}}{\underset{\text{H}}{\text{C}}}-\text{N/R}' \rightarrow ③$
- ④ Be/Al carbide $\rightarrow ①$
- ⑤ Alkene/alkyne $\rightarrow ①$


QUESTION






Identify product (A) in the following reaction:




QUESTION




QUESTION



QUESTION




~~NaI~~
H₂O Electrolysis



NaOH +
CaO, Δ



Red P
+
HI



Relation between a, b, c $\longrightarrow$ Homologues



Physical Properties



1. Heat of Combustion

$$\Delta H_{\text{combustion}} \propto \text{no. of } \underline{\text{C}}$$
$$\propto \frac{1}{\text{stability}}$$

aliphatic $\Rightarrow$ stability $\propto$ Branching

Alicyclic $\Rightarrow$ stability $\propto$ Baeyer's strain theory.

2. Physical State

$\text{C}_1 - \text{C}_4 \rightarrow$ gases

$\text{C}_4 - \text{C}_{17} \rightarrow$ liquid

$\geq \text{C}_{18} \rightarrow$ solid




$\Rightarrow$ no. of C $\uparrow$ surface area $\uparrow$ van der Waals force $\uparrow$
IMF $\uparrow$

3. Boiling Point

B. pt $\propto$ no. of 'C' atoms. (for every $\text{-CH}_2\text{-}$ increases, 20-30°C increase in B. pt)

$\propto \frac{1}{\text{Stability}}$ $\propto \frac{1}{\text{Branching}}$

B.pt order



4. Melting Point




M.pt $\propto$ no. of 'C' atoms.

$\propto$ stability $\propto$ Branching

$\propto$ symmetry.

$\Rightarrow$ For every consecutive alkanes, M.pt of even no. of 'C' alkanes $>$ M.pt of odd no. of 'C' containing alkanes



5. Density

→ alkanes are lighter than water.

→ density ↑ mass ↑

→ max. density upto 0.8 g/cm^3

6. Solubility

→ alkanes = non-polar in nature

→ soluble in non-polar solvent



Uses of Alkane:

- LPG (liquified Petroleum gas) - Cooking - Contains Propane, Butane
- LNG (liquified Natural gas) - fuel (Gasoline)
- Branching increased - Good Fuel - Example: isooctane
- CNG (Compressed Natural gas) - Vehicles fuel- contain (80-90)% Methane & traces of Ethane.
- Petrol - Vehicles fuel
- Grease (higher alkanes, hydrophobic) - used as Lubricant
- Natural gas - Methane Mainly (electricity, heat)
- Petroleum + Petrol - Dry Cleaning agent
- Kerosene oil, diesel - Fuel
- Higher Hydrocarbon - Solvent for paints






Chemical Properties of Alkanes




1. Photohalogenation of Alkanes

→ Free radical substitution



Mechanism



Types of Questions →

Monohalogenation


↓
one (H) is substituted
by one (X)





→ ① no. of different type of products.




→ ② Major/Minor.



no. of products = no. of isomers possible








Note: → Stability of F.R ⇒ $3^\circ > 2^\circ > 1^\circ$
 → Reactivity order ⇒ $F_2 > Cl_2 > Br_2 > I_2$


Fluorination




Chlorination ⇒ reactivity more }
 selectivity less }



Bromination ⇒ reactivity less }
 selectivity more }



Iodination



→ for iodination, use oxidizing agent like HNO₃ to make the rxn irreversible

Calculation of Percentage of 1°, 2° & 3°

n_1 = No. of 1° - H-atom

n_2 = No. of 2° - H- atom




n_3 = No. of 3° - H-atom

$$\% \text{ of } 1^\circ = \frac{n_1^0 R_1^0}{n_1^0 R_1^0 + n_2 R_2^0 + n_3 R_3^0} \times 100$$

$$\% \text{ of } 2^\circ = \frac{n_2^0 R_2^0}{n_1^0 R_1^0 + n_2 R_2^0 + n_3 R_3^0} \times 100$$

$$\% \text{ of } 3^\circ = \frac{n_3^0 R_3^0}{n_1^0 R_1^0 + n_2 R_2^0 + n_3 R_3^0} \times 100$$

Questions:






Combustion




a. Complete Combustion:





b. Incomplete Combustion:




4. Reaction with Steam



5. Pyrolysis/Cracking:- at high Temp / high (P), higher alkanes break into smaller molecules - alkane, alkene, H_2 .



5. Nitration




→ Free radical mechanism

→ no. of products = no. of different isomers of NO_2

→ $\boxed{\text{'C' in alkane} < 6}$ → Reactivity less
→ Reaction is done in vapour phase

→ $\boxed{C \geq 6}$ → Rxn at normal phase



6. Sulphonation



Reagent → oleum ($\text{H}_2\text{S}_2\text{O}_7$)


forming H_2SO_4 $\boxed{(\text{H}_2\text{SO}_4(\text{SO}_3))}$

→ all remaining points are same as nitration.


7. Isomerismation



➤ Catalyst → *Anhy AlCl₃ / HCl, Δ*






Branching ↑
stability ↑





8. Aromatisation







↳ Reagent $\Rightarrow$ $\text{Cr}_2\text{O}_3 / \text{Mo}_2\text{O}_3$, 15 atm.



9. Catalytic Oxidation






QUESTION



Consider the above reaction and identify the missing reagent/chemical.

[NEET-2021]



- A** B_2H_6
- B** Red phosphorus
- C** CaO ✓
- D** DIBAL-H

QUESTION



Which of the following alkane cannot be made in good yield by Wurtz reaction?


[NEET 2020]

- ☐ A 2, 3-dimethylbutane
- ☒ B n-heptane
- ☐ C n-butane
- ☐ D n-hexane

QUESTION





Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. A is [NEET 2018]



QUESTION



Which of the following reactions is expected to readily give a hydrocarbon product in good yields?
[CBSE AIPMT 1997]




QUESTION



The alkane that gives only one mono-chloro product on chlorination with Cl_2 in presence of diffused sunlight is [NEET (Odisha) 2019]

A 2, 2-dimethylbutane

B neopentane



C n-pentane

D isopentane

QUESTION



The compound that will react most readily with gaseous bromine has the formula
[NEET 2016, Phase II]

- ☐ A C_3H_6
- ☐ B C_2H_2
- ☒ C C_4H_{10}
- ☐ D C_2H_4

QUESTION



Liquid hydrocarbons can be converted to a mixture of gaseous hydrocarbons by
[CBSE AIPMT 2010]

- ☐ **A** oxidation
- ☒ **B** cracking
- ☐ **C** distillation under reduced pressure
- ☐ **D** hydrolysis

In the commercial gasolines, the type of hydrocarbons which is more desirable, is
[CBSE AIPMT 1997]

- ☒ **A** branched hydrocarbon
- ☐ **B** straight chain hydrocarbon
- ☐ **C** linear, unsaturated hydrocarbon
- ☐ **D** toluene

QUESTION



In the free radical chlorination of methane, the chain initiating step involves the formation of **[CBSE AIPMT 1994]**

- ☒ **A** chlorine free radical
- ☐ **B** hydrogen chloride
- ☐ **C** methyl radical
- ☐ **D** chloromethyl radical

QUESTION



Reactivity of hydrogen atoms attached to different carbon atoms in alkanes has the order
[CBSE AIPMT 1993]

- A** tertiary > primary > secondary
- B** Primary > secondary > tertiary
- C** Both (A) and (B)
- ~~D~~** tertiary > secondary > primary

QUESTION



Which of the following compounds has the lowest boiling point?

[CBSE AIPMT 1994]


- A** $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$
- B** $\text{CH}_3\text{CH}=\text{CH}-\text{CH}_2\text{CH}_3$
- C** $\text{CH}_3\text{CH}=\text{CH}-\text{CH}=\text{CH}_2$
- D** $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$

Alkenes





Preparation:


1. Dehydration of Alcohol:



Mechanism









Step-III → Deprotonation



→ Rate of rxn $\propto$ Stability of $\text{CH}_2 - \text{CH}_2^+$

Trick → जहाँ OH है वहाँ C^+ बनाओ, rearrange करो फिर all know बना दी।









Product
Alkene



more stable alkene
Saytzeff product




less stable alkene
Hoffmann product

QUESTION











alkyl shift







+



major

QUESTION





Which will undergoes dehydration fast?



QUESTION







Arrange according to stability of Carbocation?



QUESTION




Which amongst the following will be most readily dehydrated under acidic conditions?



2. Dehydrohalogenation of Alkyl Halides



Removal of $H+X$



product $\Rightarrow$ Saytzeff alkene (in general)
But in case of Bulky $RO^\ominus$
 $\downarrow$
Hoffmann product

$\rightarrow$ anti-elimination

$\rightarrow$ 1,2-elimination

$\rightarrow$ α, β -elimination

$\rightarrow$ E^2 mechanism


$\rightarrow$ Rate = $K [CH_3-CH_2-X] [RO^\ominus]$

$\rightarrow$ Order = molecularity = 2



$\rightarrow$ Reactivity order $\Rightarrow I^\ominus > Br^\ominus > Cl^\ominus > F^\ominus$

aqueous KOH $\rightarrow$ Nucleophile $OH^\ominus$
 $\rightarrow$ Substitution.




alcoholic KOH $\rightarrow$ Nucleophile $\Rightarrow R-O^\ominus$ (strong $Nu^\ominus$)
 $\rightarrow$ Elimination



3. Dehalogenation of Vicinal dihalides




4. From Kolbe's Electrolysis






→ free radical mechanism

5. Partial Hydrogenation of Alkyne



6. Pyrolysis/Cracking of Alkanes

→ already done



- ① Dehydration of alcohol $\rightarrow \text{C}^{\oplus}$
- ② Dehydrohalogenation of $\text{R}-\text{X}$
- ③ Dehalogenation of $\text{R}-\underset{\text{X}}{\text{CH}}-\underset{\text{X}}{\text{CH}}$
- ④ Kolbe's electrolysis
- ⑤ Alkynes
- ⑥ Pyrolysis

Physical Properties of Alkenes:



1. Heat of Hydrogenation:

always exothermic

$$\Delta H_{\text{hyd}} \propto \text{no. of } \pi\text{-bonds}$$
$$\propto \frac{1}{\text{stability}}$$

2. Physical State:

$\text{C}_2 - \text{C}_4 \rightarrow \text{gas}$
 $\text{C}_4 - \text{C}_{18} \rightarrow \text{liquid}$
 $> \text{C}_{18} \rightarrow \text{solid}$

$\rightarrow$ all are colorless/odorless
except ethene
 $\rightarrow$ pleasant smell

3. Density:


- ↳ always lower than water.
- ↳ mass ↑ density ↑
- ↳ density upto 0.8 g/cm^3 .

4. Melting Point:

- ↳ do not follow regular trend
- ↳ M.pt $\propto$ Branching $\propto$ symmetry
- $\propto$ stability
- $\propto$ Packing efficient

M.P order

Trans > Cis



5. Boiling Point:

↳ follow the regular trend.

↳ B.pt $\propto$ dipole moment

↳ In general $\Rightarrow$ cis $>$ trans

6. Solubility


↳ alkenes $\Rightarrow$ non-polar in nature.

↳ non-polar solvent में Dissolve होता है।

Chemical Properties of Alkene:



(1) Electrophilic Addition Reaction:




Symmetrical alkene




Unsymmetrical alkene





Markovnikov's rule (MKR) $\Rightarrow$ for unsymmetrical alkenes
Negative part of polar reagent attaches to that double bonded 'C' which has less no. of H attached.



Mechanism of EAR



Step-I → attack of E^+





Intermediate → Rearrangement


Step-II → Nu^- attack






Rate of EAR $\propto$ stability of C^+



Effect of Solvent







HBr (aq.)



NOTE → if solvent can provide $\text{Nu}^\ominus$, then major product will be formed by solvent $\text{Nu}^\ominus$

No






Effect of Temperature




(a) Kinetically controlled Product (KCP) $\Rightarrow$ less stable product but most stable intermediate
 $\hookrightarrow$ at low temp.



(b) Thermally controlled Product (TCP) $\Rightarrow$ more stable product irrespective of stability of intermediate
 $\hookrightarrow$ at high temp.



Step-I





Step-II



(1,2-addition)
(less stable product)
at low temp.
KCP

(1,4-addition)
(stable product)
at high temp.
TCP


Comparison of Electrophilic addition in = & -OH:




Reactivity order $\Rightarrow \underbrace{(\text{=})}_{\substack{\pi\text{-bond} \\ \text{break}}} > \underbrace{-\text{OH}}_{\substack{\sigma\text{-bond} \\ \text{break}}}$

But in case of resonance $\Rightarrow \underbrace{(\text{=})} < \textcircled{-\text{OH}}$

QUESTION



Comparison of Electrophilic addition in (=) & (≡)





(Vinyl carbocation) $\rightarrow$ less stable


Reactivity for EAR $\Rightarrow (=) > (\equiv)$

except in case of resonance


QUESTION



Stereochemistry of electrophilic addition reaction:







⇒ if chiral centre is present, equal amount of (d) & (l).



QUESTION



Alkene $\xrightarrow{\text{HCl}}$ Enantiomeric pair or Racemic mixture which of the following is possible Alkene?




9. Non-Classical Carbocation:



→ No rearrangement

→ partial (+ve) charge on 'C'



→ used as test for unsaturation

Br₂/CCl₄ → colored.

When attaches on (=) bond, become colorless.

→ anti-addition

→ stereochemistry

→ In case of unsymmetrical alkene → follow Markovnikov's rule.

CAT / CAR

Cis $\xrightarrow{\text{Anti addition}}$ Racemic mixture / Threo.

TAE / TAM



Trans $\xrightarrow{\text{anti addition}}$ Meso / Erythro

Cis $\xrightarrow{\text{syn addition}}$ Meso / Erythro. (CSM/CSE)


Trans $\xrightarrow{\text{syn addition}}$ Racemic mixture / Threo (TSR/TST)

Mechanism



Step-1 (Addition of Electrophile)




→ 3 membered
cyclic Transition
state



Step-2 (Addition of Nucleophile)



Effect of solvent. (NCC)



Reactant	Reagent	E ⁺	Nu ⁻ (major)	Major Product
CH ₃ - CH = CH ₂	Br ₂ / <u>ROH</u>	Br ₂	Br [⊖] / RO [⊖]	$\begin{array}{c} \text{Br} \\ \\ \text{CH}_3 - \underset{\text{OR}}{\text{CH}} - \text{CH}_2 \end{array}$
Ph - CH = CH ₂	Cl ₂ / <u>PhOH</u>	Cl ₂	Cl [⊖] / PhO [⊖]	$\begin{array}{c} \text{Cl} \\ \\ \text{CH}_3 - \underset{\text{OPh}}{\text{CH}} - \text{CH}_2 \end{array}$
CH ₃ - CH = CH ₂	Br - Cl	Br-Cl	Cl [⊖]	$\begin{array}{c} \text{Br} \\ \\ \text{CH}_3 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_2 \end{array}$
Ph - CH = CH ₂	I - Cl	ICl	Cl [⊖]	$\begin{array}{c} \text{I} \\ \\ \text{Ph} - \underset{\text{Cl}}{\text{CH}} - \text{CH}_2 \end{array}$
CH ₃ - CH = CH ₂	HO - Cl	HOCl	OH [⊖]	$\begin{array}{c} \text{Cl} \\ \\ \text{CH}_3 - \underset{\text{OH}}{\text{CH}} - \text{CH}_2 \end{array}$
Ph - CH = CH ₂	HO - I	HOI	OH [⊖]	$\begin{array}{c} \text{I} \\ \\ \text{Ph} - \underset{\text{OH}}{\text{CH}} - \text{CH}_2 \end{array}$

QUESTION


What will be the product?






CAR
TAM

- ☐ A Racemic Mixture
- ☒ B Meso compound
- ☐ C Threo
- ☐ D Both (A) & (C)

(a) Reaction with cold conc. H_2SO_4



(b) Reaction with water in presence of acid



(c) Antimarkoni/Peroxide/Kharasch Effect





→ Negative part of polar reagent will attach to that double bonded 'C' which has more no. of 'H' attached.

→ free radical mechanism.

→ No rearrangement

(HBr in presence of $R-O-O-R$ Catalyst)
Peroxide

all steps are exothermic




$\left\{ \begin{array}{l} HF \\ HCl \end{array} \right\} \rightarrow$ all steps are endothermic
 $\left(\begin{array}{l} H-I \end{array} \right) \rightarrow$ first step exo then endo

Mechanism
initiation







Propagation



Termination





2. Oxymercuration & Demercuration of Alkenes:





→ Reagent $\Rightarrow$ (i) $\text{Hg}(\text{OAc})_2 / \text{H}_2\text{O} - \text{THF}$

(ii) NaBH_4

→ anti-addition.

→ stereochemistry $\rightarrow$ CAR/CAT
TAM/TAE

→ product will be formed $\Rightarrow$ Markovnikov's rule without rearrangement.




(i) $\text{Hg}(\text{OAc})_2 / \text{H}_2\text{O} / \text{THF}$
(ii) NaBH_4



Trick

-OH $\rightarrow$ from H_2O

-H $\rightarrow$ from NaBH_4






3. Hydroboration - Oxidation of Alkene:

→ Reagent $\Rightarrow$ (i) B_2H_6 or $(BH_3)_2$
(ii) H_2O_2/OH^-

→ water is added by Anti-Markovnikov's rule

Trick

$H \rightarrow BH_3/B_2H_6$
 $OH \rightarrow H_2O_2$



4. Hydroxylation of Alkene:

→ Product → vicinal diol will be formed.

→ Reagent →

syn-addition

CSM/CSE


TSR/TST





(i) cold dilute alkaline KMnO_4 (Bayer's reagent)
(1%)

or

(ii) OsO_4 / pyridine


+ $\text{NaHSO}_3 / \text{H}_2\text{O}$





5. Hydrogenation of Alkenes:

→ syn-addition










6. Oxidation of alkenes:



a. by Hot Alk. KMnO_4 (strong oxidising agent) OR Acidic KMnO_4 OR

Acidic $\text{K}_2\text{Cr}_2\text{O}_7$







b. By Ozone (Ozonolysis)










⇒ Oxidative Ozonolysis → Reagent → $\boxed{O_3}$
↳ product same as ~~hot~~ $KMnO_4$ / acidic $KMnO_4$ / $K_2Cr_2O_7$.








⇒ Reductive Ozonolysis ⇒ Reagent ① O_3
② Zn/H_2O or Pb/H_2O or $(CH_3)_2S$













NOTE → ozonolysis is preferred for aliphatic than aromatic

7. Addition of Oxygen:


Reagent: Ag_2O or per-acid ($\text{R}-\overset{\text{O}}{\parallel}\text{C}-\text{O}-\text{O}-\text{H}$)






8. Alkylation:



9. Polymerisation:





QUESTION



10. Substitution in Alkenes:

↳ allylic substitution




Reagent $\Rightarrow \text{Br}_2 / 773\text{K}$

or
 $\text{Cl}_2 / 773\text{K}$

or
NBS (N-Bromosuccinimide)
773K
↳ Best for $\text{Br}^\bullet$.

free radical
mechanism.









11. Reaction of Alkene with KMnO_4 and NaIO_4 :



$\text{KMnO}_4/\text{NaIO}_4$ → same as oxidative ozonolysis.

$\text{KMnO}_4/\text{NaIO}_4$
 ↳ Lemux reagent



QUESTION




The major product formed in dehydrohalogenation reaction of 2-bromo pentane is pent-2-ene. This product formation is based on [NEET 2021]

☒ **A** Saytzeff's rule

☐ **B** Hund's rule

☐ **C** Hoffmann rule




☐ **D** Huckel's rule



QUESTION



2, 3-dimethyl-2-butene can be prepared by heating which of the following compounds with a strong acid: [NEET 2016, P-I]



QUESTION





When 3, 3-dimethyl-2-butanol is heated with conc. H_2SO_4 , the major product obtained is
[CBSE AIPMT 1995]

☒ A 2, 3-dimethyl-2-butene

☐ B 3, 3-dimethyl-2-butene

☐ C 2, 3-dimethyl-1-butene

☐ D cis and trans-isomers of 2, 3-dimethyl-2-butene







QUESTION






An alkene of ozonolysis gives methanal as one of the product. Its structure is

[NEET Sep.-2020]







QUESTION








An alkene 'A' on reaction with O_3 and $\text{Zn} - \text{H}_2\text{O}$ gives propanone and ethanal in equimolar ratio. Addition of HCl to alkene 'A' gives 'B' as the major product. The structure of product 'B' is

[NEET (National) 2019]



QUESTION



QUESTION




Which of the following compounds with molecular formula, C_5H_{10} yields acetone on ozonolysis?
[CBSE AIPMT 2007]

☒ A 2-methyl-2-butene

☐ B 3-methyl-1-butene

☐ C Cyclopentane





☐ D 2-methyl-1-butene



QUESTION



Which one of the following alkenes will react faster with H_2 under catalytic hydrogenation conditions? (R = alkyl substituent) [CBSE AIPMT-2005]



QUESTION



Reaction of HBr with propene in the presence of peroxide gives


[CBSE AIPMT 2004]

A iso-propyl bromide

B 3-bromo propane

C allyl bromide

D n-propyl bromide



QUESTION



The alkene $R - CH = CH_2$ reacts readily with B_2H_6 and formed the product B which on oxidation with alkaline H_2O_2 produces [CBSE AIPMT 1995]




- ☐ A $R - CH_2 - CHO$
- ☒ B $R - CH_2 - CH_2 - OH$
- ☐ C $R - C \begin{matrix} \text{= O} \\ | \\ CH_3 \end{matrix}$
- ☐ D $R - CH \begin{matrix} | \\ OH \end{matrix} - CH_2 \begin{matrix} | \\ OH \end{matrix}$

QUESTION



In the reaction with HCl, an alkene reacts in accordance with the Markownikoff's rule, to give a product 1-chloro-1methylcyclohexane. The possible alkane is

[CBSE AIPMT 2015]



QUESTION



Among the following alkenes

1-butene

cis-2-butene

trans-2-butene

I

II


III

the decreasing order of stability is

[CBSE AIPMT 2000]

A $II > I > III$

☒ B $III > II > I$



C $III > I > II$


D $I > II > III$

QUESTION



Compound X on reaction with O_3 followed by Zn/H_2O gives formaldehyde and 2-methyl propanal as products. The compound X is: [NEET 2022]

- ☐ A Pent-2-ene
- ☒ B 3-Methylbut-1-ene
- ☐ C 2-Methylbut-1-ene
- ☐ D 2-Methylbut-2-ene



QUESTION



The compound, $\text{CH}_3 - \overset{\text{CH}_3}{\underset{|}{\text{C}}} = \text{CH} - \text{CH}_3$

on reaction with NaOI_4 , in the presence of cold KMnO_4 gives [CBSE AIPMT 2003]

- A** $\text{CH}_3\text{COCH}_3 + \text{CH}_3\text{CHO}$
- B** $\text{CH}_3\text{CHO} + \text{CO}_2$
- C** CH_3COCH_3
- D** $\text{CH}_3\text{COCH}_3 + \text{CH}_3\text{COOH}$





Alkynes

Introduction:





➤ S-Character → s-character ↑ acidic nature ↑ Electronegativity ↑



➤ Type of Alkynes:

→ Terminal alkyne





→ non-terminal alkyne



Preparation:


1. Dehydrohalogenation of Vicinal Dihalides





2. Dehydrohalogenation of Geminal Dihalides



3. Dehydrohalogenation of Vicinal Tetrahalides



4. Dehalogenation of Haloform (CHX_3)





→ Symmetrical alkynes will be formed

5. Higher Alkynes from Lower Alkynes




↳ having acidic (H)
↳ Terminal alkyne




6. Preparation of Acetylene only ($\text{CH} \equiv \text{CH}$)





(a) From Kolbe's Electrolysis



(b) From Graphite




(c) From Calcium Carbide (Industrial Scale)






Al Carbide $\rightarrow \text{CH}_4$
Be Carbide $\rightarrow \text{CH}_4$



Preparation of Calcium Carbide:



QUESTION



QUESTION



Physical Properties of Alkynes:



1. Physical state

$C_2 - C_4 \rightarrow$ gas

$C_5 - C_{12} \rightarrow$ liquid

$> C_{12} \rightarrow$ solid

colorless / odorless

except



if pure $\rightarrow$ pleasant smell

if impure $\rightarrow$ garlic smell

2. Density $\rightarrow$ lighter than water.

max $\uparrow$ density $\uparrow$

max upto 0.69 g/cm^3

3. Melting Point/Boiling point

- ↓ alkynes have higher m.pt/b.pt than alkenes/alkanes
- M.pt $\propto$ symmetry/stability
- B.pt high due to π -bonds


4. Solubility

- non-polar in nature
- soluble in non-polar solvent



Chemical Properties of Alkynes:




➤ Electrophilic Addition Reaction



(a) Addition of Hydrogen (Hydrogenation)





(b) Addition of Halogen (Halogenation)





(MKR)



(c) Addition of Halogen acids (Hydrohalogenation)




e) Addition of Hypohalous Acid (HOX)






f) Addition of Water in presence of Acid






phenyl > methyl > ethyl ketone
ketone




Addition of Hydrogen Cyanide (HCN):




Addition of acetic acid:




Addition of Arsenic Trichloride:





2. Nucleophilic Addition Reaction:



EAR





NAR




3. Oxidative - Hydroboration (HBO):

$\xrightarrow{\text{H}_2\text{O}}$ addition
 $\xrightarrow{\text{Reagent} \Rightarrow \text{B}_2\text{H}_6/\text{H}_2\text{O}_2/\text{OH}^-}$
 $\xrightarrow{\text{Antimarkovnikov}}$




4. Acetylide Formation:

(a) Reaction with Ag/Cu:





(b) Reaction with Na:



(c) Reaction with Grignard Reagent:




(d) Reaction with Tollen's Reagent:



5. Oxidation of Alkynes:





(a) Oxidation in Neutral KMnO_4 :

Similar to Reductive Ozonolysis.






(b) Oxidation with cold dil. Alkaline KMnO_4

→ in case of alkene → hydroxylation




(c) Oxidation By Hot/Acidic KMnO₄ → same as Oxidative Ozonolysis.








6. Ozonolysis:

(a) Reductive Ozonolysis (Similar to Neutral KMnO_4)






(b) Oxidative Ozonolysis (Similar to Hot/ H^+ KMnO_4)






7. Polymerisation Reaction of Alkynes:


(a) Cyclic Polymerisation





(b) Vinylic Polymerisation



c) Linear Polymerisation:





QUESTION


$R-CH_2-CCl_2-R \xrightarrow{\text{Reagent}} R-C \equiv C-R$. The reagent is

- A** Na
- B** HCl in H_2O
- C** KOH in C_2H_5OH ✓
- D** Zn in alcohol

QUESTION



The most suitable reagent for the following conversion, is:



[NEET (National) 2019]

- A** H_2 , Pd/C, quinoline ✓✓
- B** Zn/HCl
- C** $\text{Hg}^{2+}/\text{H}^+$, H_2O
- D** Na/liquid NH_3

QUESTION



Which one is the correct order of acidity?

[NEET 2017]

- A** $\text{CH}_2 = \text{CH}_2 > \text{CH}_3 - \text{CH} = \text{CH}_2 > \text{CH}_3 - \text{CH} \equiv \text{CH} > \text{CH} \equiv \text{CH}$
- B** $\text{CH} \equiv \text{CH} > \text{CH}_3 - \text{C} \equiv \text{CH} > \text{CH}_2 = \text{CH}_2 > \text{CH}_3 - \text{CH}_3$
- C** $\text{CH} \equiv \text{CH} > \text{CH}_2 = \text{CH}_2 > \text{CH}_3 - \text{C} \equiv \text{CH} > \text{CH}_3 - \text{CH}_3$
- D** $\text{CH}_3 - \text{CH}_3 > \text{CH}_2 = \text{CH}_2 > \text{CH}_3 - \text{C} \equiv \text{CH} > \text{CH} \equiv \text{CH}$


QUESTION



In the reaction,



[NEET 2016, P-I]



X and Y are

- A** X = 2-butyne; Y = 3-hexyne
- B** X = 2-butyne; Y = 2-hexyne
- C** X = 1-butyne; Y = 2-hexyne
- D** X = 1-butyne; Y = 3-hexyne

QUESTION




Which of the following reagents will be able to distinguish between 1-butyne and 2-butyne?
[CBSE AIPMT 2012]

☒ **A** NaNH_2

☐ **B** HCl

☐ **C** O_2

☐ **D** Br_2




QUESTION



When acetylene is passed through dil. H_2SO_4 in presence of HgSO_4 the compound formed is
[CBSE AIPMT 1999]

- A** ether
- B** ketone
- C** acetic acid
- D** acetaldehyde ✓



QUESTION



The cylindrical shape of an alkyne is due to

[CBSE AIPMT 1997]

- ☐ A three sigma C-C bonds
- ☐ B three π C - C bonds
- ☐ C two sigma C - C and one π C - C bonds
- ☒ D one sigma C - C and two π C - C bonds



QUESTION



A compound is treated with NaNH_2 to give sodium salt. Identify the compound.

[CBSE AIPMT 1993]



QUESTION



Reduction of 2-butyne with sodium in liquid ammonia gives predominantly
[CBSE AIPMT 1993]







- ☐ A cis-2-butene
- ☒ B trans-2-butene
- ☐ C no reaction
- ☐ D n-butane



QUESTION



Which is the most suitable reagent among the following to distinguish compound (III) from rest of the compounds? [CBSE AIPMT 1989]



Acetylenic hydrogens are acidic because

[CBSE AIPMT 1989]

- ☒ **A** sigma electron density of C-H bond in acetylene is nearer to carbon which has 50% s-character
- ☐ **B** acetylene has only one hydrogen on each carbon
- ☐ **C** acetylene contains least number of hydrogens among the possible hydrocarbons having two carbons
- ☐ **D** acetylene belong to the class of alkynes with molecular formula C_nH_{2n-2}

Benzene

Benzene




Introduction:



Toluene



ortho/para activating for E^+




Aromatic Compounds $\rightarrow$ Benzenoid
 $\rightarrow$ non-benzenoid


Preparation:



1. Decarboxylation of Sodium Benzoate




2. From Chlorobenzene




3. From Phenol




4. From Sulphonic Acid



5. Synthesis/Reforming/Aromatisation




6. From Benzene Diazonium Chloride




7. Manufacture



a. n-Hexane (Reforming/Catalytic Reforming/Aromatisation)




b. Coal Tar



Physical Properties of Benzene:

- Physical State: *liquid*
- M.P./B.P.: $\propto$ *no. of C*
- Nature: *non-polar*
- Solubility: *soluble in non-polar solvent*
- Aromatic Compounds : *producing pleasant smell.*
- On Burning:
 - Sooty flame
 - ↳ *because of high 'C' content*

Chemical Properties of Benzene:




1. Aromatic Electrophilic Substitution Reaction:




Mechanism

Step-I








Step-II








$FeCl_3 / FeBr_3$
 Lewis acid














Can undergo Rearrangement



Friedel craft acylation








Benzoylation





Acetylation

except $R-\overset{\overset{O}{\parallel}}{C}^{\oplus}$ do not
 have $3^{\circ}C$ in R
 if R has $3^{\circ}C$ → then that
 C will be carbocation




+




Note:






Friedel craft Reaction in Aryl & Vinyl Carbocation.




Other ways to generate E⁺/R⁺ (Alkylation)



Intramolecular Friedel Craft Reaction: 5/6 membered ring




Halogenation in the present of dark:



→ Electrophilic substitution rxn

→ Hexachlorobenzene

halogenation in the presence of sunlight:




Benzenehexachloride (BHC)

→ Gammaxane


→ insecticide
(lindane)

C.I. = 8


Hydrogenation:




Combustion of Benzene:



Ozonolysis:




Dehydrogenation:




Oxidation:

(a) Controlled oxidation in Air





oxidation of Benzene
↳ not by oxidising agent
↳ Under drastic condition.

Oxidation of Alkyl Benzene:




CH_3
 CH_3
 C
 CH_3
 CH_3
 + maleic + CO_2 + H_2O
 anhydride



QUESTION



Some meta-directing substituents in aromatic substitution are given. Which one is most deactivating?
[NEET 2013]



QUESTION



Which of the following compounds will not undergoes Friedel-Craft's reaction easily?
[NEET 2013]

- ☐ A Cumene
- ☐ B Xylene
- ☒ C Nitrobenzene
- ☐ D Toluene

QUESTION



Nitrobenzene can be prepared from benzene by using a mixture of conc. HNO_3 and conc. H_2SO_4 . In the mixture, nitric acid acts as a/an **[CBSE AIPMT 2009]**

- A** reducing agent
- B** acid
- C** base
- D** catalyst

QUESTION



Benzene reacts with CH_3Cl in the presence of anhy. AlCl_3 to form

[CBSE AIPMT 2009]

- ☒ **A** toluene
- ☐ **B** chlorobenzene
- ☐ **C** benzylchloride
- ☐ **D** xylene

QUESTION



Which one of these, is not compatible with arenes?

[CBSE AIPMT 1998]

- ☐ **A** Greater stability
- ☐ **B** Delocalisation of π -electrons
- ☒ **C** Electrophilic additions
- ☐ **D** Resonance


QUESTION



Benzene reacts with n-propyl chloride in the presence of anhydrous AlCl_3 to give
[CBSE AIPMT 1993]



- A** 3-propyl-1-chlorobenzene
- B** n-propyl benzene X
- C** no reaction
- D** iso-propyl benzene ✓





QUESTION



Which one of the following is most reactive towards electrophile.


[CBSE AIPMT 2008]



QUESTION



Among the following, the reaction that proceeds through an electrophilic substitution, is
[NEET (National) 2019]






QUESTION



Which one of the following is a free radical substitution reaction?

[CBSE AIPMT 2003]



QUESTION




Among the following compounds (I-III), the correct order of reaction with electrophile is
[CBSE AIPMT 1997]

A $\text{II} > \text{III} > \text{I}$

B $\text{III} < \text{I} < \text{II}$

C $\text{I} > \text{II} > \text{III}$


D $\text{I} > \text{III} > \text{II}$



I




II



III



Summary



———— FOR NOTES & DPP CHECK DESCRIPTION ————



Homework

→ Selection



———— **FOR NOTES & DPP CHECK DESCRIPTION** ————

THANK YOU

