PRACHAND NEEL




ONE SHOT ®



Botany

Cell: The Unit of Life

Rupesh Chaudhary Sir





Topics tope covered

1 Cell: The Unit of Life




जिन घरों में <mark>खाने के लिए रोटी</mark> नहीं होती, वहाँ घरों में Motivational Video नहीं देखा जाता ! करना है तो करना है।




By- Rupesh Sir

- Detail Study
- 2. Flow chart / Diagrammatic Based
- Covered Complete NCERT
- All PYQ
- Notes (Short Notes in Detail)
- 6. 5 Hours Duration
- 7. 14 Years का Experience है Paid Batch के Equivalent Content दूंगा.
- 8. After Class, Comment करके बताना Lecture कैसा लगा।

9) आपने मेरे NOTES पड़ लिये तो NCERT भी पड़ने का NEED नहीं है मेने सारा NOTES NCERT को DECODE करके तैयार किया है LET'S TAKE A CHALLENGE






MATTHIAS SCHLEIDEN, 1838
GERMAN BOTANIST

STUDY: ALL PLANTS

COMPOSED OF CELLS
WHICH FORMS TISSUE



THEODORE SCHWANN, 1839
BRITISCH ZOOLOGIST

* STUDY: ANIMAL CELL --> COVERED BY THIN LAYER
PLASMA MEMBRANE

* STUDY: PLANT CELL -- CELL WALL PRESENSE: ONIQUE CHARACTER

HYPOTHESIS: PLANTS & ANIMALS

COMPOSED


CELL & ITS PRODUCT (TISSUE)


NEW CELL ARISE BY DIVISION OF PRE EXISTING CELL (1)
Omnis Cellula e Cellula





2) ALL LIVING ORGANISM: COMPOSED: CELL & PRODUCT OF CELL


OVERVIEW OF CELL : EUKARYOTE CELL. >ONION PEEL: PLANT : CELL WALL → DELIMITING LAYER > HUMAN CHEEK : ANIMAL : CELLMEMBRANE] NUCLEOPLASM ds linear dna, → HISTONE PROTEIN (BASIC) NON HISTONE PROTEIN (ACIDIC) RNA




















REDUCED/DIMINISHED IN , CAPABLEOF (add of cellulose)
MATURECELL GROWTH CONTINUENCY OUTER, YOUNG CELL, Confinuously THIN

INNER, MATURE CELL, THICK. (NEAR TO CELL MEMBRANE)

> CONNECT CYTOPLASM OF TWO CEUS. CELLWALL (PW+SW), MIDDLE LAMELLA CROSS/TRAVERSED BY PLAS MODESMATA.


> > FUNCTION

- SHAPE
- * PROTEIN PRESENT IN CW PROTECT FROM PATHOGEN/INFECTION
- PROVIDE BARRIER TO UNDESIRABLE MACROMOLECULE (LARGE SIZE)
- CELL TO CELL INTERACTION.

COMMON LAYER B/W TWO CELL /

(CALCIUM PECTATE)

- > OUTER TO CELL MEMBRANE (MINERAL)
- AIGAE: CELLULOSE, GALACTAN, MANNAN, COCO3
- > PLANT: CELLULOSE, HEMICELLULOSE, PECTIN, PROTEIN
- 3 FUNGI : CHITIN



P.P. PERIPHERAL PROTEIN, EASY


I.P. INTEGRAL PROTEIN, DIFFICULT

TOTALY PARTIALLY (I LAYER)

BURIED BURIED

(2 LAYER)

COMMON LIPID: PHOSPHOLIPID





FCHEMICAL STUDY: HUMAN RBC: CELL MEMBRANE (UPID+PROTEIN)


& BIOCHEMICAL STUDY


Membrane: Caubohydrate also present

RBC Membrane 52% P THIS RATIO
40% L VARY IN

Main Funct
(TRANPORT)

DIFFERENT
CELL MEMBRANE





H ~


LOW CONL


MOVE
THROUGH SELECTIVELY PERMEABLE /
CELL MEMBRANE
(OSMOSIS)

ACTIVE TRANSPORT

- → LOW HIGH CON" (SOME MOLECULES/
- ⇒ AGAINST CONCENTRATION
- = ATP/ENERGY
- 4 CARRIER PROTEIN

eg. Nat Kt pump





RIBOSOME

PROTEIN GYNTHESIS, MEMBRANLESS
SMALLEST ORGANELLE, PALADE PARTICLE.

```
PROK.
RIBONUCLEOPROT- - 705 (RNA+PROTEIN)
    EIN
                           505 (LARGE)
       30s (small)
                       > 55, 2357RNA
      6SYRNA
                       = 34 PROTEIN
      21 PROTEIN
                   EUK)
                   809.
          405
                              605.
    7 18SYRNA
                           = 155, 285, 5.85
      33 P.
                   80 Protein
```

```
S: Svedbergy ONIT


IS SEDIMENTATION (VELOCITY)

COEFFICIENT HIGH

HIGH

(Indirect measure

Size & density)
```



TUBUUNI PROTEIN


PERIPHERAL MICROTUBULE: (M.T)

=) 9 (doublet)

a I doubles: 2MTCA

7) 9X2 = 18 M·T

CENTRE: 2 MT (Parrof MT)



(Peaphery centre. (doublet) + 29x2=18+2 720 M.T.

⇒Central M·T Connected BY BRIDGE


> Central M.T Covered By central SHEATH

= RADIAL SPOKE: 9


arise from one of TWO (A MT)

> INTERDOUBLET/LINKERS CONNECT TWO PERIPHERAL DOUBLET

CENTROSOME & CENTRIQUE




- CENTROSOME: 2 CENTRIOLE CYLINDRICAL
 - STRUCTURE)
- 90 to each other
- => membranless, cout wheel Structure





PROTENAEOUS

STRUCTURE

PORMADION







- () Vesicle come from ER (PROTEIN & LIPID)
- 2) moves to cis side of GB
- 3) P&L enters into vesicle of GB.
- A) PRL moves to cisternae.

- 5 some protein move outside the cell (secretary protein)
- Some protein after modification Release inside the cell in form of Lysosome
- D Lysosome formed due to packaging
 By 9.B. But RER provide

 protein for synthesis of Lysosome


ENDOMEMBRANE SYSTEM: ER, GB, LYSOSOME, VACUOLE (WORK IN COORDINATION)

NON ENDOMEMBRANE SYSTEM: MITOCHONDRIA, (NOT WORK IN COORDINATION)

PEROXISOME

vsasome formation by G.B. + membrane FOOD DIGESTION Carbohydrase 7 acidic professe - DIGESTIVE/ medium Lipase HYDROLYTIC PH: 5. ENZYME Núclease

VACUOLE PLANTCELL, LARGE, OCCUPY 90/ VOLUME OF CELL SAP.




PRACHAND SERIES


TELEGRAM CHANNEL








LYSOSOME














NOCLEUS

- > ROBERT BROWN, 1831
- 7 Cell: generally: one nucleus.
- 7 paramecium: Two nucleus.
- => Mature sleve Tube, Thucleus mammalian RBC ABSENT LIVING
 - (1) NUCLEAR MEMBRANE
 - & DOUBLE LAYER.
 - 7 PERINUCLEAR SPOCE/ NM SEPERATE CYTOPLASMIP NUCLEOPLASM
 - = protein & RNA: BIDIRECTIONAL MOVEMENT THROUGH NUCLEAR PORE.





Chromosome






Figure 8.1 Dtagram showing different shapes of the cells




Figure 8.2 Diagram showing comparison of eukaryotic cell with other organisms



HEET

7. Cytoskeleton



An elaborate network of filamentous proteinaceous structures consisting of microtubules, microfilaments and intermediate filaments present in the cytoplasm is collectively referred to as the cytoskeleton. The cytoskeleton in a cell are involved in many functions such as mechanical support, motility, maintenance of the shape of the cell.

myosin

NCICLEAR LAMINA (INNER NUCLEAR

WEMBEANT)





Many membrane bound minute vesicles called microbodies that contain various enzymes, are present in both plant and animal cells.



Match List I with List II: (2024)

Choose the correct answer from the options given below:

- 1 A-IV, B-III, C-II, D-I
- 2 A-IV, B-II, C-III, D-I
- 3 A-II, B-IV, C-I, D-III
- A-II, B-I, C-IV, D-III

	List I		List II
A.	Axoneme	j.	Centriole
В.	Cartwheel pattern	all.	Cilia and flagella
C.	Crista	iii.	Chromosome
D.	Satellite	IV.	Mitochondria



The DNA present in chloroplast is:

- 1 Linear, double stranded
- 2 Circular, double stranded
- 3 Linear, single stranded
- 4 Circular, single stranded

(2024)



Match List I with List II: (2024)

Choose the correct answer from the options given below:

- 1 A-IV, B-III, C-II, D-I
- 2 A-IV, B-II, C-III, D-I
- 3 A-II, B-IV, C-I, D-III
- 4 A-II, B-I, C-IV, D-III

	List-I	ĕ	List-II
A.	Nucleolus 1.		Site of formation of glycolipid
В.	Centriole II		Organization like the cartwheel
C.	Leucoplasts	I.	Site for active ribosomal RNA synthesis
D.	Golgi apparatus IV	V.	For storing nutrients



Movement and accumulation of ions across a membrane against their concentration gradient can be explained by (2023)

- 1 Facilitated Diffusion
- 2 Passive Transport
- 3 Active Transport
- 4 Osmosis



How many different proteins does the ribosome consist of?



- **1**) 60
- **2** 40
- 3 20
- 4 80

(2023)



Which of the following are NOT considered as the part of endomembrane system? (2023)

A. Mitochondria 🗸

B. Endoplasmic reticulum

C. Chloroplasts <

D. Golgi complex

E. Peroxisomes

Choose the most appropriate answer from the options given below:

- 1 A, C and E only
- 2 A and D only
- 3 A, D and E only
- 4 B and D only



Which of the following functions is carried out by cytoskeleton in a cell?

(2023)

- 1 Protein synthesis
- 2 Motility
- 3 Transportation
- 4 Nuclear division



Given below are two statements:

(2023 manipur)

Statement-I: In bacteria, the mesosomes are formed by the extensions of plasma membrane.



Statement-II: The mesosomes, in bacteria, help in DNA replication and cell wall formation. In the light of the above statements, choose the most appropriate answer from the options given below:

- 1 Statement-I is correct but Statement-II is incorrect.
- 2 Statement-I is incorrect but Statement-II is correct.
- 3 Both Statement-I and Statement-II are correct.
- 4 Both Statement-I and Statement-II are incorrect.



Which of the following statements are correct with respect of Golgi apparatus?

A. It is the important site of formation of glycoprotein and glycolipids. (2023 manipur)

B. It produces cellular energy in the form of ATP.

C. It modifies the protein synthesized by ribosomes on ER.

D. It facilitates the transport of ions.

E. It provides mechanical support.

Choose the most appropriate answer from the options given below:

- 1 B and C only
- 2 A and C only
- 3 A and D only
- 4 D and E only



Match the List-I with List-II.

Choose the correct answer from the options given below:

D-II A-II B-1. C-IV (2022)

- 1 A-Q, B-P, C-S, D-R
- 2 A-S, B-R, C-Q, D-P
- 3 A-P, B-R, C-S, D-Q
- 4 A-R, B-Q, C-P, D-S

	List-I		List-II
A.	Metacentric chromosome	l.	Centromere situated close to the end forming one extremely short and one very long arms
B.	Acrocentric chromosome	II.	Centromere at the terminal end
C.	Submetacentric	III.	Centromere in the middle forming two equal arms of chromosomes
D.	Telocentric chromosome	IV.	Centromere slightly away from the middle forming one shorter arm and one longer arm



Which of the following statements with respect to Endoplasmic Reticulum is incorrect? (2022)

- SER are the sites for lipid synthesis
- 2 RER has ribosomes attached to ER
- 3 SER is devoid of ribosomes
- In prokaryotes only RER are present

 ER absent)



Which type of substance would face difficulty to pass through the cell membrane?

2022 (Phase 2)

- POLAR INATERSOURIE Substance with hydrophobic moiety
- Substance with hydrophilic moiety
- All substance irrespective of hydrophobic and hydrophilic moiety
- Substance soluble in lipids



If the pH in lysosomes is increased to alkaline, what will be the outcome? 2022 (Phase 2)

- 1 Hydrolytic enzymes will function more efficiently
- 2 Hydrolytic enzymes will become inactive
- 3 Lysosomal enzymes will be released into the cytoplasm
- 4 Lysosomal enzymes will be more active



Which of the following is an incorrect statement?

(2021)

- 1) Microbodies are present both in plant and animal cells.
- The perinuclear space forms a barrier between the materials present inside the nucleus and that of the cytoplasm.
- Nuclear pores act as passages for proteins and RNA molecules in both directions between nucleus and cytoplasm.
- Mature sieve tube elements possess a conspicuous nucleus and usual cytoplasmic organelles.



Match the List-I with List-II Choose the correct answer from the options given below.

(A)-(I); (B)-(IV); (C)-(III); (D)-(II)

(A)-(III); (B)-(IV); (C)-(I); (D)-(II)

(A)-(II); (B)-(III); (C)-(IV); (D)-(I)

(A)-(IV); (B)-(III); (C)-(II); (D)-(I)

	List-I		List-II
A.	Cristae	I.	Primary constriction in chromosome
В.	Thylakoids	II.	Disc-shaped sacs in Golgi apparatus
C.	Centromere	1 II.	Infoldings in mitochondria
D.	Cisternae	IV.	Flattened membranous sacs in stroma of plastids

(2021)



When the centromere is situated in the middle of two equal arms of chromosomes, the chromosome is referred as:

(2021)

- 1 Telocentric
- 2 Sub-metacentric
- 3 Acrocentric
- 4 Metacentric



The organelles that are included in the endomembrane system are:

(2021)

- 1 Endoplasmic reticulum, Golgi complex, Lysosomes and Vacuoles.
- 2 Golgi complex, Mitochondria, Ribosomes and Lysosomes.
- 3 Golgi complex, Endoplasmic reticulum, Mitochondria and Lysosomes.
- 4 Endoplasmic reticulum, Mitochondria, Ribosomes and Lysosomes.



Which of the following statements about inclusion bodies is incorrect?

(2020)

- 1 These are involved in ingestion of food particles.
- 2 They lie free in the cytoplasm
- 3 These represent reserve material in cytoplasm C
- 4 They are not bound by any membrane C



Which is the important site of formation of glycoproteins and glycolipids in eukaryotic cells? (2020)

- 1 Peroxisomes
- 2 Golgi bodies
- 3 Polysomes
- 4 Endoplasmic reticulum



The biosynthesis of ribosomal RNA occurs in:

- 1 Golgi apparatus
- 2 Microbodies
- 3 Nucleolus V
- 4 Ribosomes

(2020 Covid)



Inclusion bodies of blue-green, purple and green photosynthetic bacteria are:(2020 Covid)

- (1) Gas vacuoles ~
- 2 Centrioles
- 3 Microtubules
- 4 Contractile vacuoles



Match the following Lists and select the correct option;

(2020 Covid)

	(A)	(B)	(C)	(D)
	(III)	(1)	(11)	(IV)
2	(IV)	(11)	(1)	(III)
3	(1)	(11)	(III)	(IV)
4		(1)		(IV)

	List-I		List-II
A.	Smooth Endoplasmic Reticulum	ı. Y	Protein synthesis
B.	Rough endoplasmic reticulum	II.	Lipid synthesis
C.	Golgi complex	/ III.	Glycosylation
D.	Centriole	IV.	Spindle formation



The size of Pleuropneumonia - like Organism (PPLO) is:

- 1 2 μm
- **2** 10 20 μm
- **3** 0.1 μm
- **4** 0.02 μm

(2020 Covid)



Which of the following pair of organelles does not contain DNA?

- 1 Mitochondria and Lysosomes
- 2 Chloroplast and Vacuoles
- 3 Lysosomes and Vacuoles
- 4 Nuclear envelope and Mitochondria

(2019)



The shorter and longer arms of a sub-metacentric chromosome are referred to as (2019)

- s-arm and l-arm respectively
- 2 p-arm and q-arm respectively
- q-arm and p-arm respectively
- m-arm and n-arm respectively



Which of the following statements is not correct?

(2019)

- 1 Lysosomes have numerous hydrolytic enzymes.
- 2 The hydrolytic enzymes of lysosomes are active under acidic pH.
- 3 Lysosomes are membrane bound structures.
- 4 Lysosomes are formed by the process of packaging in the endoplasmic reticulum.



The concept of "Omnis cellula" regarding cell division was first proposed by (2019)

- Rudolf Virchow
- 2 Theodor Schwann
- 3 Schleiden
- 4 Aristotle



Which of the following statements regarding mitochondria is incorrect?

(2019)

- Outer membrane is permeable to monomers of carbohydrates, fats and proteins.
- 2) Enzymes of electron transport are embedded in outer membrane.
- 3 Inner membrane is convoluted with infoldings.
- 4) Mitochondrial matrix contains single circular DNA molecule and ribosomes.



Which of the following cell organelles is present in the highest number in secretory cells?

1) Mitochondria (2019 odisha)

2 Golgi complex

SECRETION (proken all outsid):

Secreting proken)

packaging modification:

delivered

3 Endoplasmic reticulum

4 Lysosomes



Non-membranous nucleoplasmic structures in nucleus are the site for active synthesis of (2019 odisha)

- 1 protein synthesis
- 2 mRNA
- 3 rRNA
- 4 tRNA





Which of the following nucleic acids is present in an organism having 70 S ribosomes only?

(2019 odisha)

- 1) Single stranded DNA with protein coat
- 2 Double stranded circular naked DNA
- 3 Double stranded DNA enclosed in nuclear membrane
- 4 Double stranded circular DNA with histone proteins



Match the List-I with List-II.

(2019 odisha)

Choose the right match from options given below:

- 1 A-R, B-S, C-Q, D-P
- 2 A-S, B-R, C-P, D-Q
- 3 A-R, B-Q, C-S, D-P
- 4 A-P, B-Q, C-S, D-R

	List-I		List-II
A.	Golgi apparatus	P.	Synthesis of protein
В.	Lysosomes products	Q.	Trap waste and excretory
C.	Vacuoles	R.	Formation of glycoproteins and glycolipids
D.	Ribosomes	S.	Digesting biomolecules



Which among the following is not a prokaryote?

- 1) Saccharomyces (yeast) Eukanyola ~
- 2) Mycobacterium
- 3 Nostoc
- 4 Oscillatoria



Many ribosomes may associate with a single mRNA to form multiple copies of a polypeptide simultaneously. Such strings of ribosomes are termed as

- 1 Polysome
- 2 Polyhedral bodies
- 3 Plastidome
- 4 Nucleosome



The Golgi complex participates in

- 1 Fatty acid breakdown
- 2 Formation of secretory vesicles
- Respiration in bacteria
- 4 Activation of amino acid



Which of the following events does not occur in rough endoplasmic reticulum?

- 1 Protein folding
- 2 Protein glycosylation
- 3 Cleavage of signal peptide
- 4 Phospholipid synthesis



Which of the following is true for nucleolus?

- 1 Larger nucleoli are present in dividing cells.
- 2 It is a membrane-bound structure.
- 3 It takes part in spindle formation.
- 4 It is a site for active ribosomal RNA synthesis



[OS] (2017-Delhi)

Which of the following components provides sticky character to the bacterial cell?

- 1 Cell wall
- 2 Nuclear membrane
- 3) Plasma membrane
- 4 Glycocalyx



Which of the following cell organelles is responsible for extracting energy from carbohydrates to form ATP? (2017-Delhi)

- 1 Lysosome
- 2 Ribosome
- 3 Chloroplast
- 4 Mitochondrion



A complex of ribosomes attached to a single strand of RNA is known:

(2016 - I)

- 1 Polysome
- 2 Polymer
- 3 Polypeptide
- 4 Okazaki fragment



Mitochondria and chloroplast are

- Semi-autonomous organelles
- B. Formed by division of pre-existing organelles and they contain DNA but lack protein synthesizing machinery

Which one of the following options is correct?

(2016-I)

- 1) Both (A) and (B) are correct
- (B) is true but (A) is false
- (A) is true but (B) is false
- Both (A) and (B) are false



Microtubules are the constituents of:

- Cilia, Flagella and Peroxisomes
- 2 Spindle fibres, Centrioles and Cilia
- 3 Centrioles, Spindle fibres and Chromatin
- 4 Centrosome, Nucleosome and Centrioles

(2016-I)



Select the wrong statement:

(2016 - II)

- 1) Cyanobacteria lack flagellated cells.
- 2 Mycoplasma is a wall-less microorganism
- 3 Bacterial cell wall is made up of peptidoglycan.
- Pilli and fimbriae are mainly involved in motility of bacterial cells



Select the mismatch:

- 1 Protists-Eukaryotes
- 2 Methanogens-Prokaryotes
- 3 Gas vacuoles-Green bacteria
- 4 Large central vacuoles-Animal cells

(2016 - II)



A cell organelle containing hydrolytic enzymes is:

(2016 - II)

- 1 Ribosome
- 2 Mesosome
- 3 Lysosome
- 4 Microsome

