# SEAL

# DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

**Test Booklet Series** 

T. B. C.: PGT - 4/21



#### **TEST BOOKLET**

PAPER - II BIOLOGY

40585

SI. No.

Time Allowed: 2 Hours

Maximum Marks : 100

#### : INSTRUCTIONS TO CANDIDATES :

- IMMEDIATELY AFTER THE COMMENCEMENT OF THE EXAMINATION, YOU SHOULD CHECK THAT THIS TEST BOOKLET DOES NOT HAVE ANY UNPRINTED OR TORN OR MISSING PAGES OR ITEMS ETC. IF SO, GET IT REPLACED BY A COMPLETE TEST BOOKLET OF THE SAME SERIES ISSUED TO YOU.
- ENCODE CLEARLY THE TEST BOOKLET SERIES A, B, C OR D, AS THE CASE MAY BE, IN THE APPROPRIATE PLACE IN THE ANSWER SHEET USING BALL POINT PEN (BLUE OR BLACK).
- You have to enter your Roll No. on the Test Booklet in the Box provided alongside. DO NOT write anything else on the Test Booklet.
- 4. YOU ARE REQUIRED TO FILL UP & DARKEN ROLL NO., TEST BOOKLET / QUESTION BOOKLET SERIES IN THE ANSWER SHEET AS WELL AS FILL UP TEST BOOKLET / QUESTION BOOKLET SERIES AND SERIAL NO. AND ANSWER SHEET SERIAL NO. IN THE ATTENDANCE SHEET CAREFULLY, WRONGLY FILLED UP ANSWER SHEETS ARE LIABLE FOR REJECTION AT THE RISK OF THE CANDIDATE.
- 5. This Test Booklet contains 100 items (questions). Each item (question) comprises four responses (answers). You have to select the correct response (answer) which you want to mark (darken) on the Answer Sheet. In case, you feel that there is more than one correct response (answer), you should mark (darken) the response (answer) which you consider the best. In any case, choose ONLY ONE response (answer) for each item (question).
- You have to mark (darken) all your responses (answers) ONLY on the separate Answer Sheet provided, by using BALL POINT PEN (BLUE OR BLACK). See instructions in the Answer Sheet.
- 7. All items (questions) carry equal marks. All items (questions) are compulsory. Your total marks will depend only on the number of correct responses (answers) marked by you in the Answer Sheet. There will be no negative markings for wrong answers.
- Before you proceed to mark (darken) in the Answer Sheet the responses (answers) to various items (questions) in the Test Booklet, you have to fill in some particulars in the Answer Sheet as per the instructions sent to you with your Admission Certificate.
- After you have completed filling in all your responses (answers) on the Answer Sheet and after conclusion of the examination, you should hand over to the Invigilator the Answer Sheet issued to you. You are allowed to take with you the candidate's copy / second page of the Answer Sheet along with the Test Booklet, after completion of the examination, for your reference.
- 10. Sheets for rough work are appended in the Test Booklet at the end.

DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

- Algal partner in Lichens can be:
  (B)
  - (A) Cyanobionts
  - (B) Phycobionts
  - (C) Cyanobionts and phycobionts
  - (D) Rhodophyta
- Specialized hyphal structures for nutrient uptake by parasitic fungus is :
  - (A) Arbuscules
  - (B) Haustoria
  - (C) Mycelium
  - (D) Rhizomorphs
- 90 percent of vascular plants show mycorrhizal association except plants from:
  - (A) Amranthaceae
  - (B) Chenopodiaceae
  - (C) Brassicaceae
  - (D) All of these
- Aflatoxins, a highly carcinogenic liver toxin is produced by:
  - (A) Claviceps
  - (B) Amanita
  - (C) Candida
  - (D) Aspergillus
- In Bryophytes, the 'Alternation of generation' involves :
  - (A) Haploid gametophyte alternates with diploid sporophyte

- (B) Diploid gametophyte alternates with haploid sporophyte
- (C) Haploid gametophyte alternates with haploid sporophyte
- (D) Diploid gametophyte alternates with diploid sporophyte
- 6. In Lycopodium, plant body is differentiated into:
  - (A) Adventitious root, stem,rhizophores and leaves
  - (B) Underground rhizome, rhizophores and leaves
  - (C) Stem, rhizoids and scaly leaves
  - (D) Root, stem, rhizoids and leaves
- 7. Prothallus is:
  - (A) Zygote developing into sporophyte
  - (B) Zygote developing into gametophyte
  - (C) Spores developing into an independent gametophyte
  - (D) Spores developing into an independent sporophyte

- 8. Gymnosperms appeared during:
  - (A) Cretaceous
  - (B) Jurassic
  - (C) Triassic
  - (D) Permian
- 9. The 'Maiden fern' tree is:
  - (A) Marselia
  - (B) Selaginella
  - (C) Cycas
  - (D) Ginkgo biloba
- Vascular phanerogams are :
  - (A) Bryophytes
  - (B) Pteridophytes
  - (C) Gymnosperms
  - (D) Angiosperms
- 11. Following is true regarding the Crustaceans:
  - (A) Metanauplius is the first fundamental stage in all crustaceans
  - (B) Nauplius larva is the first fundamental stage in all crustaceans
  - (C) Protozoea is the first fundamental stage in all crustaceans
  - (D) Zoea is the first fundamental stage in all crustaceans

- 12. Microscopic Molluscs are:
  - (A) Monoplacophorans
  - (B) Aplacophorans
  - (C) Polyplacophorans
  - (D) Sacphopoda
- 13. Echinoderms are:
  - (A) Protostomes with exoskeleton
  - (B) Deuterostomes with endoskeleton
  - (C) Protostomes with radial symmetry
  - (D) Protostomes with watervascular system
- 14. One of the following is not a basic chordate character:
  - (A) Notochord
  - (B) Ventral nerve chord
  - (C) Pharyngeal gill slits
  - (D) Post-anal tail
- 15. A unique set of cells that originate above the neural tube during development is:
  - (A) Neural arch
  - (B) Centrum
  - (C) Neural crest cells
  - (D) Neural tube

- 16. First fishes known were jawless with heavy bony armor are usually referred to as:
  - (A) Agnathans
  - (B) Chondrichthyes
  - (C) Osteichthyes
  - (D) Placoderms
- 17. The first terrestrial vertebrates were :
  - (A) Tetrapods
  - (B) Lancelet
  - (C) Pisces
  - (D) None of these
- 18. Kinetic skull is a feature of :
  - (A) Lizard
  - (B) Snakes
  - (C) Both (A) and (B)
  - (D) None of these
- Archaeopteryx was considred a bird because of :
  - (A) Feathers on wings and tail
  - (B) Solid bones
  - (C) Lack of breast bone
  - (D) Long reptilian tail
- 20. Egg laying mammals are:
  - (A) Marsupials
  - (B) Monotremes

- (C) Placentals
- (D) None of these
- 21. Cell number is controlled by:
  - (A) Cell division
  - (B) Apoptosis
  - (C) Both cell division and apoptosis
  - (D) None of these
- 22. Lampbrush chromosomes are formed during the :
  - (A) Leptotene stage of meiotic prophase I
  - (B) Zygotene stage of meiotic prophase I
  - (C) Pachytene stage of meiotic prophase I
  - (D) Diplotene stage of meiotic prophase I
- 23. Polytene chromosomes are formed as a result of :
  - (A) Repeated rounds of DNA replication
  - (B) Lack of cell division
  - (C) Repeated DNA replication without cell division
  - (D) Repeated DNA replication with cell division

MJ-4A/20 (4)

- 24. Signal peptides have a common characteristic tripartite structure composed of:
  - (A) A positively charged N-terminus (n region)
  - (B) Stretch of 7-15 hydrophobic residues (h region)
  - (C) A polar region that often contains helix-breaking proline and glycine residues as well as the signal peptide cleavage site (c region)
  - (D) All of these

#### 25. GAAP:

- (A) Resides in the Mitochondria and protects cells from apoptosis
- (B) Resides in the Endoplasmic reticulum and protects cells from apoptosis
- (C) Resides in the Golgi and protects cells from apoptosis
- (D) Resides in the Nucleus and protects cells from apoptosis
- 26. Mitochondrial disorder due to mutation in mitochondrial DNA (mtDNA) is:
  - (A) Keams-Sayre syndrome

- (B) Leber's hereditary optic neuropathy
- (C) Pearson's syndrome
- (D) All of these
- 27. Prokaryotes have 70S ribosomes are composed of:
  - (A) Small (20S) and a large (50S) subunits
  - (B) Small (30S) and a large (40S) subunits
  - (C) Small (30S) and a large (50S) subunits
  - (D) Small (10S) and a large (60S) subunits
- 28. Eukaryotic cytoskeleton is composed of :
  - (A) Microfilaments
  - (B) Intermediate filaments
  - (C) Microtubules
  - (D) Microfilaments, intermediate filaments and microtubules
- 29. Lamin proteins is part of:
  - (A) Nuclear lamina
  - (B) Nucleolus
  - (C) Nuclear pore
  - (D) Nuclear membrane

- Endoplasmic reticulum forms the interconnected network of :
  - (A) Tubules
  - (B) Vesicles
  - (C) Cisternae
  - (D) Tubules, vesicles and cistemae
- 31. The core activators of the cell cycle control system is / are:
  - (A) Cyclin Dependent Kinases
  - (B) Cyclin proteins
  - (C) Kinases
  - (D) Phosphorylase
- 32. Brakes on the cell cycle are provided by:
  - (A) Rb protein
  - (B) p16, p21
  - (C) p27
  - (D) All of these
- 33. The LOD score (logarithm (base 10) of odds) is a statistical test used for:
  - (A) Crossing-over studies
  - (B) Linkage analysis
  - (C) Crossing-over and linkage
  - (D) None of these
- 34. Haemophilia A and B are:
  - (A) X-linked genetic disorders
  - (B) Autosomal dominant disorders
  - (C) Autosomal recessive disorders
  - (D) Non-genetic disorders

- 35. Mutation in 'Dystrophin' gene leads to:
  - (A) Duchenne Muscular Distrophy (DMD)
  - (B) Becker Muscular Distrophy (BMD)
  - (C) Duchenne Muscular Distrophy (DMD) and Becker Muscular Distrophy (BMD)
  - (D) Neurofibromatosis
- 36. Cystic fibrosis has following symptoms:
  - (A) Muscle wasting
  - (B) Dementia
  - (C) Abnormal bleeding
  - (D) Progressive lung damage
- Polygenic inheritance in human can be exemplified by :
  - (A) Height
  - (B) Eye colour
  - (C) Hair colour
  - (D) All of these
- 38. 1:2:1 is the:
  - (A) The phenotypic ratio of the incomplete dominance
  - (B) The genotypic ratio of the incomplete dominance
  - (C) The genotypic and the phenotypic ratio of the incomplete dominance
  - (D) The genotypic ratio of Lethality

- 39. Deviation from the Mendel's dihybrid cross is shown by :
  - (A) Epistasis
  - (B) Hypostasis
  - (C) Complementation
  - (D) All of these
- 40. Polyploid cells in humans occur in highly differentiated cells such as :
  - (A) Liver parenchyma
  - (B) Heart muscle
  - (C) Bone marrow
  - (D) All of these
- 41. A multitude of protein structures are rapidly determined by :
  - (A) X-ray crystallography
  - (B) Nuclear Magnetic Resonance (NMR)
  - (C) Circular Dichroism spectroscopy (CD)
  - (D) All of these
- 42. α-helices, β-sheets and turns are defined by :
  - (A) Torsion angles between backbone atoms  $C_{i-\ell} N_i Ci\alpha C_i$
  - (B)  $N_i Ci\alpha C_i N_{i+\ell}$
  - (C) Regular main chain hydrogen bonding patterns
  - (D) All of these

- 43. Aromatic amino acid are:
  - (A) Phenylalanine, Tyrosine, Tryptophan
  - (B) Glycine, Tyrosine, Tryptophan
  - (C) Phenylalanine, Serine, Tryptophan
  - (D) Phenylalanine, Tyrosine, Methionine
- 44. Amino acids with amide side-chains are :
  - (A) Glycine, Glutamine
  - (B) Asparagine, Glutamine
  - (C) Asparagine, Alanine
  - (D) Valine, Glutamine
- 45. In sickle-cell anemia:
  - (A) Valine mutates to Histidine
  - (B) Valine mutates to Proline
  - (C) Glutamic acid mutates to valine
  - (D) Glutamic acid mutates to Leucine
- The Michaelis-Menten constant (K<sub>m</sub>) estimates :
  - (A) Substrate concentration required for an enzyme to reach one-half its maximum reaction rate
  - (B) Enzyme concentration required to reach one-half its maximum reaction rate
  - (C) Number of substrate molecules handled by one active site per second
  - (D) Number of substrate molecules handled by several active site per second

#### 47. An irreversible inhibitor:

- (A) Binds to an allosteric site
- (B) Binds to an Enzyme-substrate complex
- (C) Permanently inactivates the enzymes
- (D) Strongly resembles the actual substrate

#### 48. Carbonic anhydrase:

- (A) Uses Zn as a cofactor bound to its active site
- (B) Uses Zn as a cofactor bound to its allosteric site
- (C) Uses Cu as a cofactor bound to its active site
- (D) Uses Cu as a cofactor bound to its allosteric site

#### 49. Gluconeogenesis:

- (A) Generates glucose using carbohydrate source
- (B) Generates glucose using noncarbohydrate carbon substrates
- (C) Generates lipids using carbohydrate source
- (D) Generates lipid using noncarbohydrate carbon substrates

- 50. Fatty acids are metabolically oxidized at their:
  - (A) α-carbon atom
  - (B) β-carbon atom
  - (C) y-carbon atom
  - (D) λ-carbon atom

#### 51. The variant forms of DNA are:

- (A) A, C, D, E, Z
- (B) F, C, D, E, Z
- (C) A, G, D, E, Z
- (D) A, C, H, E, Z
- 52. Two types of gene groups / clusters are :
  - (A) Operons
  - (B) Multigene families
  - (C) Operons and Multigene families
  - (D) Pseudogene

#### 53. Open Reading Frame (ORF) is:

- (A) A set of codons that runs discontinuously and is bound by initiation codon at start and termination codon at the end
- (B) A set of codons that runs continuously and is bound by initiation codon at start and termination codon at the end
- (C) A set of codons that runs continuously and is bound by initiation codon at both the end
- (D) A set of codons that runs discontinuously and is bound by termination codon at both the end

- 54. In E. coli termination occurs at:
  - (A) Promoter site
  - (B) Enhancer site
  - (C) Palindromes
  - (D) Silencer site
- 55. Third base degeneracy or Wobbling serves to minimize the effect of :
  - (A) Mutation
  - (B) DNA Repair
  - (C) Proof-reading function
  - (D) None of these
- 56. During translation:
  - (A) GTP is used to charge tRNA and in removing secondary structure from mRNA while ATP is used for ribosome movement and binding of accessory factors while
  - (B) GTP is used for ribosome movement and binding of accessory factors while ATP is used to charge tRNA and in removing secondary structure from mRNA
  - (C) GTP and ATP are used for ribosome movement and binding of accessory factors
  - (D) GTP and ATP are used to charge tRNA and in removing secondary structure from mRNA

- 57. Separation of double helix during replication is achieved by :
  - (A) Single strand binding proteins
  - (B) Ligase
  - (C) Helicase enzyme
  - (D) Primase
- 58. DNA binding domains in eukaryotes are :
  - (A) Helix-tum-helix
  - (B) Zinc fingers
  - (C) Basic domains
  - (D) Helix-turn-helix, Zinc fingers and Basic domains
- 59. Chromosomal scaffold that is involved in the binding with chromatin fibres are largely composed of:
  - (A) Acidic non-histone nuclear protein Topoisomerase II
  - (B) Histone nuclear protein Topoisomerase II
  - (C) Both histone and acidic nonhistone nuclear protein Topoisomerase II
  - (D) Acidic non-histone cytoplasmic protein Topoisomerase II
- 60. 5-bromouracil (5-BU) is a base analog derived from:
  - (A) Adenine
  - (B) Guanine
  - (C) Cytosine
  - (D) Thymine

61. Major group of blood plasma (C) Alveoli proteins are: (D) Pharynx (A) Albumin Pyelonephritis is a urinary tract 66. (B) Globulins infection caused by: (C) Fibrinogen (A) Bacteria (D) Albumin, Globulins and (B) Virus Fibrinogens (C) Protozoa Opening in the fetal heart in the 62. (D) Fungi interatrial septum is: 67. The functional and structural unit of Fossa ovalis (A) the kidney is: (B) Foramen ovale Bowman's capsule (A) (C) Septum primum (D) Semilunar valves (B) Glomerulus (C) Nephron 63. Cardiac Output (CO) can be (D) Loop of Henle estimated by Stroke Volume (SV) and Heart Rate (HR) by an equation: 68. Major endocrine feedback systems (A) CO = SV + HR that are mediated through hypo-(B) CO = SV - HR thalamus-pituitary are: (C) CO = SV × HR (A) TRH-TSH-T3/T4 (D) CO = SV + HR (B) CRH-ACTH-Cortisol Renin-Angiotensin-64. Obligate nasal breathers are : (C) Aldosterone (A) Human All of these (D) (B) Horse (C) Birds 69. The beginning of gastrulation is (D) Elephant marked by: (A) Embryonic disc 65. Respiratory gaseous exchange takes place at: (B) Primitive streak

(B)

(A) Trachea

Bronchi

(C)

Blastocoel

(D) Cleavage furrow

70. The functional k	kidney in human is:
----------------------	---------------------

- (A) Pronephros
- (B) Mesonephros
- (C) Metanephros
- (D) All of these
- 71. Water potential (ψ<sub>w</sub>) of a plant cell is given by inter-relationship between solute potential or osmotic potential (ψ<sub>s</sub>) and pressure potential or hydrostatic pressure (ψ<sub>p</sub>) through a equation:
  - (A)  $\psi_w = \psi_s + \psi_p$
  - (B)  $\psi_w = \psi_s \psi_p$
  - (C)  $\psi_w = \psi_s \times \psi_p$
  - (D)  $\psi_w = \psi_s + \psi_p$

# 72. Loss of water in plants is regulated by:

- (A) Change in the vascular diameter
- (B) Trichomes
- (C) Leaf cuticle
- (D) All of these

#### 73. Photosynthesis occurs via:

- (A) Light dependent reactions to synthesize NADPH and ATP
- (B) Light independent reactions to capture and reduce CO<sub>2</sub>
- (C) Both (A) and (B)
- (D) None of these

- 74. P<sub>680</sub> and P<sub>700</sub> are the two special molecules of:
  - (A) Chlorophyll A
  - (B) Chlorophyll B
  - (C) Carotenes
  - (D) Phycobilins

## 75. Cryptochrome mediated photoresponses are :

- (A) Phototropism
- (B) Stomatal opening
- (C) Phototaxis
- (D) All of these

#### 76. Skotomorphogenesis is:

- (A) Regulatory effect of light on plant cell, tissue growth
- (B) Regulatory effect of light on the differentiation of plant cell, tissue
- (C) Development of seedling in dark
- (D) None of these

## During seed germination the level of Abscisic acid (ABA) :

- (A) Increases
- (B) Decreases
- (C) Remains the same
- (D) None of these

- 78. Secondary xylem differentiation is promoted by :
  - (A) Auxin
  - (B) Cytokinin
  - (C) Ethylene
  - (D) Gibberellin
- 79. Of the following the gaseous phytohormone is:
  - (A) Salicyclic acid
  - (B) Strigolactones
  - (C) Ethylene
  - (D) Jasmonates
- Test-tube fertilization of angiosperms was invented by :
  - (A) B. M. John
  - (B) P. Maheswari
  - (C) W. A. Jensen
  - (D) S. G. Nawaschin
- 81. Lymphocytes and antibodies constitute:
  - (A) First line of defence
  - (B) Second line of defence
  - (C) Third line of defence
  - (D) None of these
- Antibody dependent cell mediated toxicity (ADCC) :
  - (A) Used to destroy large organisms that cannot be phagocytosed

- (B) Used to destroy virus
- (C) Used to destroy bacteria
- (D) Natural killer cells are not involved in the process
- 83. Capsules, cell walls, toxins, viral capsids, and flagella etc, constitute:
  - (A) Antigen of microbial nature
  - (B) Antigen of non-microbial nature
  - (C) Antigen of both microbial and non-microbial nature
  - (D) Haptens
- 84. Autoimmune disorders can be exemplified by:
  - (A) Systemic Lupus Erythematosus
  - (B) Rheumatoid Arthritis
  - (C) Hashimoto's Thyroiditis
  - (D) All of these
- 85. Major criteria for bacteriophage classification is :
  - (A) Phage morphology
  - (B) Nucleic acid properties
  - (C) Phage morphology and nucleic acid properties
  - (D) None of these

86. Temperate phages used : 91. Density-dependent limiting factors of a population are: (A) Lysogenic cycles Lytic cycles (A) Intrapopulation competition (B) Lysogenic and lytic cycles (C) (B) Predation (D) None of these (C) Disease and parasites Lambda phage (λ) parasitizes : (D) All of these (A) Escherichia coli 92. Prey-predator interactions show their Clostridium botulinum (B) numbers oscillate over time (C) Vibrio cholera depicting: (D) Streptococcus pyogenes (A) Sigmoid curve Enterovirus follows: 88. (B) J-shaped curve (A) Respiratory transmission (C) Bell shaped curve (B) Faecal-oral transmission (D) Wave shaped curve (C) Sexual transmission 93. Human and other large mammal (D) Blood borne transmission populations are: Generation of novel Influenza A 89 (A) r-selected viruses is due to: (B) k-selected (A) Antigenic drift (C) Both r- and k-selected (B) Antigenic shift (D) None of these (C) Antigenic drift and antigenic shift Simpson dominance index for a 94. (D) Altered cytokine expression stream inhabiting 55 percent of blackfly larvae and 9 other species, Cervical carcinoma and benign warts 90. 5 percent each will be: are caused by: (A) 3.25 (A) Epstein Barr virus (B) 0.325 (B) Human Papilloma virus

(C) Hepatitis C virus

(D) Human T cell Leukaemia virus

(C) 32.5

(D) 325

- 95. Species diversity depends on:
  - (A) Species richness
  - (B) Species evenness
  - (C) Both species richness and evenness
  - (D) None of these
- 96. Herbivores (primary consumers) constitute the :
  - (A) First trophic level
  - (B) Second trophic level
  - (C) Third trophic level
  - (D) None of these
- 97. Various types of food chain in an ecosystem are:
  - (A) Grazing food chain
  - (B) Saprophytic / Detritus food chain
  - (C) Parasitic food chain
  - (D) All of these

- 98. Percentage flow of energy across various trophic levels is:
  - (A) 10 percent
  - (B) 20 percent
  - (C) 30 percent
  - (D) 40 percent
- 99. Functioning of an ecosystem depends on:
  - (A) Energy flow
  - (B) Material cycling
  - (C) Energy flow and material cycling
  - (D) None of these
- 100. Major gaseous pollutants are:
  - (A) Carbon monoxide
  - (B) Sulphur dioxide
  - (C) Chlorofluorocarbons
  - (D) All of these

# SPACE FOR ROUGH WORK

MJ-4A/20 (15) PGT-4/21

mary sold and the second of th